arXiv:1304.0834v1 [math.AP] 3 Apr 2013

Long-time asymptotics for the porous medium equation: The spectrum of the linearized operator

Christian Seis*

April 4, 2013

Abstract

We compute the complete spectrum of the displacement Hessian operator $\mathcal{H}\psi = -m\rho_*^{m-2}\nabla\cdot(\rho_*\nabla\psi)$, which is obtained from the confined porous medium equation in dimensions $N \geq 2$ by linearization along vector fields $\nabla\psi$ around its stationary attractor ρ_*. Here, ρ_* denotes the Barenblatt profile and ψ is an element in the weighted Sobolev space $H^1_{\rho_*}$ that arises in a natural way as the tangent plane in the (formal) gradient flow interpretation of the porous medium equation for the entropy $E(\rho)$. On a formal level, the operator $\mathcal{H}$ is conjugate to the Hessian of $E(\rho)$ via similarity transformation. We show that $\mathcal{H}$ can be understood as a self-adjoint operator on $H^1_{\rho_*}$ and find that its spectrum is purely discrete. The knowledge of the complete spectrum and the explicit information about the corresponding eigenfunctions give new insights on the convergence and higher order asymptotics of solutions to the porous medium equation towards its attractor ρ_*. More precisely, the inspection of the eigenfunctions allows to identify symmetries in $\mathbb{R}^N$ with flows whose rates of convergence are faster than the uniform, translation-governed bound. The present work imitates the analogous study of Denzler & McCann [9] for the fast-diffusion equation and complements their results in the porous medium regime.

1 Introduction

In this paper, we study the long-time asymptotics of nonnegative solutions to the porous medium equation, i.e.,

$$\partial_t\rho - \Delta(\rho^m) \;=\; 0 \quad \text{in } \mathbb{R}^N, \tag{1}$$

with exponent $m > 1$. This equation is best known for modeling the flow of an isentropic gas through a porous medium; other applications include groundwater infiltration, population dynamics, and heat radiation in plasmas (cf. [31, Ch. 2]).

*Department of Mathematics, University of Toronto, 40 St. George Street, M5S 2E4, Toronto, Ontario, Canada

Solutions to (1) feature different phenomena depending on the degree of the nonlinearity ρ^m. In the case $m = 1$, equation (1) is the ordinary diffusion (or heat) equation. For $0 < m < 1$, the diffusion flux $m\rho^{m-1}$ diverges as ρ vanishes and thus, for suitable initial data, the solution spreads over the whole $\mathbb{R}^N$ immediately. In this situation, equation (1) is often called the *fast diffusion* equation. In the porous-medium range $m > 1$, the diffusion flux increases with the density and degenerates where $\rho = 0$. Consequently, solutions will preserve a compact support and hence this type of propagation goes by the name *slow diffusion*. In the present paper, we restrict our attention exclusively to the latter case. For a study of the fast diffusion equation and more general evolutions of porous-medium type, we refer to VÁZQUEZ [30] and references therein.

The Cauchy problem for the porous medium equation is solved in various settings. As the equation is degenerate parabolic, that is, it is parabolic only where the solution is positive, solutions are in general not classical. More precisely, if the initial datum is zero in some open subset of $\mathbb{R}^N$, then there is a slowly propagating free boundary that separates the region where the solution is positive from the region where it is zero. For suitable initial configurations, e.g. $0 \le \rho_0 \in L^1(\mathbb{R}^N)$, unique strong solutions are known to exists, and these solutions are bounded and continuous. Moreover, strong solutions preserve total mass, $\|\rho(t,\,\cdot\,)\|_{L^1(\mathbb{R}^N)} = \|\rho_0\|_{L^1(\mathbb{R}^N)}$ for all $t > 0$. The Cauchy problem for the porous medium equation is reviewed by VÁZQUEZ in [31].

It is well-known [35, 3, 24] that the porous medium equation allows for self-similar solutions, so-called Barenblatt solutions, propagating on the scale $|x| = t^\alpha$ where

$$\alpha \;=\; \frac{1}{N(m-1)+2},$$

and given by

$$\rho_*(t,x) \;=\; \frac{1}{t^{N\alpha}}\left(L - \frac{\alpha(m-1)}{2m}\frac{|x|^2}{t^{2\alpha}}\right)_+^{1/(m-1)}. \tag{2}$$

Here, $(\cdot)_+ = \max\{\,\cdot\,, 0\}$ and L is an arbitrary constant that can be fixed, for instance, by normalizing the total mass or by choosing the radius of the support of ρ_*. It is interesting to note that although the Barenblatt solution is a strong solution of the porous medium equation, it is not a solution of the Cauchy problem as ρ_* diverges to the delta measure δ_0 (times a constant depending on L) at time zero — a reason for which it is often called a "source-type solution".

The Barenblatt solution describes the long-time behavior of any solution ρ having same total mass as ρ_*. Indeed, for arbitrary non-negative initial data in L^1, solutions spread with a finite propagation speed over the whole space and, while diffusing, the shape of the solution becomes asymptotically close to the profile of the Barenblatt solution,

$$\rho(t,\,\cdot\,) \approx \rho_*(t,\,\cdot\,) \quad \text{as } t \gg 1.$$

This long-time behavior was intensively studied over many years, starting with the work by KAMIN [15, 16], who established uniform convergence in one space dimension. The generalization to several dimensions is due to FRIEDMAN & KAMIN [12] and KAMIN & VÁZQUEZ [17]. These authors use similarity (rescaling) transformations and identify the limit orbits for the porous medium equation via compactness arguments. A completely different approach relies on energy (or entropy) methods, a way prepared by RALSTON in [27], who established a first convergence result in L^1 in one dimension. This approach was generalized independently by CARRILLO & TOSCANI, OTTO, and DEL PINO & DOLBEAULT in [6, 22, 7] to arbitrary dimensions. In these articles, instead of using geometric properties of the density function, the authors study the abstract energy landscape and compute the decay rates of the energy (or entropy) functional. In [6] and [22], these decay rates are converted into bounds on the asymptotics in L^1 and in the Wasserstein distance, respectively. We refer to VÁZQUEZ' excellent review article [29] for an almost complete discussion of the long-time asymptotics of solutions to the porous medium equation, including a discussion on the optimal rates of convergence.

The investigation of the optimal rate at which solutions converge to the Barenblatt solution is tied together with the choice of the initial data. The minimal requirement of integrability of the initial data — which is from the physical point of view natural when thinking of ρ_0 as a concentration density — is mathematically necessary as solutions to non-integrable initial data, i.e., $\rho_0 \in L^1_{\mathrm{loc}} \setminus L^1$, show a different behavior in the long-time limit (cf. [29, Cor. 1.2]). The authors in [6, 22, 7] derive algebraic rates of convergence for finite entropy solutions, cf. (5) below. The exponent in the algebraic rate is estimated by α, which is optimal for spatial translations of the Barenblatt solution. The goal of the present article is to go one step further and to study the complete spectrum of a linearized version of the porous medium equation. The knowledge of the spectrum and of all eigenfunctions not only yields information about the slowest converging modes (like a spectral gap estimate would do), but also allows to extract information about the characteristic geometric pathologies to all orders. In particular, we will study the role of translations, shears, and dilations for the convergence rates of solutions towards its attractor ρ_*. While spectral properties of the heat equation (i.e., (1) with $m = 1$) are well-know in any space dimension, the investigation of the spectrum of the porous medium equation is completely new in the multi-dimension case. In the one-dimensional setting, ANGENENT [2] computed the long-time asymptotics to all orders for solutions with compactly supported initial data. An earlier formal spectral analysis is due to ZEL'DOVIC & BARENBLATT [34]. Progress in the analogous problem for the fast diffusion equation was recently obtained by DENZLER, KOCH & MCCANN [8]: the authors compute higher order asymptotics in weighted Hölder spaces, building up on earlier results for the linearized fast diffusion equation derived by DENZLER & MCCANN [9].

Our approach to compute the spectrum of the linearized porous medium equation follows closely the work of Denzler & McCann [9], who studied the analogous prob-

lem for the fast diffusion equation. Their work is based on a method which is fairly common in the quantum mechanics literature, more precisely in the spectral analysis of Schrödinger operators, see e.g. [21]. The main difference between the analysis in [9] and the one in the present work relies on the occurrence of a free boundary in the porous medium regime. In particular, opposed to the situation for $m \leq 1$, the Barenblatt solution has compact support in $\mathbb{R}^N$. This fact will play a crucial role in the definition of the linearized operator as ("asymptotic") boundary conditions on the boundary of the Barenblatt solution have to be taken into consideration. Somehow surprisingly (and unexpected, see [9, p. 303]), compared to the analog study in [9] for the fast diffusion regime, the analysis of our linearized operator simplifies thanks to the compact support of ρ_* in many aspects.

We caution the reader that our linearization of the porous medium equation performed in subsection 1.3 is only formally justified. In particular, the rigorous results obtained for the linearized operator have to be considered as conjectures for the asymptotics of solutions to the nonlinear equation (1).

1.1 Rescaling

In order to investigate the asymptotic behavior of solutions to the porous medium equation, it is convenient to rescale variables in such a way that the Barenblatt solution becomes a stationary solution, that means $|x| \sim t^\alpha$ in view of (2). One convenient choice of new variables is

$$x = \beta t^\alpha \hat{x}, \quad t = \exp(\alpha^{-1}\hat{t}\,), \quad \text{and} \quad \rho = (\alpha\beta^2)^{1/(m-1)} t^{-\alpha N}\hat{\rho},$$

with

$$\beta = \left(\frac{2mL}{\alpha(m-1)}\right)^{1/2},$$

to the effect of

$$\rho(t,x) = \frac{(\alpha\beta^2)^{1/(m-1)}}{t^{\alpha N}}\hat{\rho}\left(\alpha \ln t, \frac{x}{\beta t^\alpha}\right).$$

We remark that the logarithmic rescaling of the time variable, $\hat{t} = \alpha \ln t$, guarantees the parabolic structure of the equation. In fact, with its new variables, equation (1) turns into the nonlinear Fokker–Planck equation

$$\partial_{\hat{t}}\hat{\rho} - \hat{\nabla}\cdot(\hat{x}\hat{\rho}) - \hat{\Delta}(\hat{\rho}^m) \;=\; 0, \tag{3}$$

sometimes also called the *confined* porous medium equation. Moreover, with that particular choice of β, the rescaled, time-independent Barenblatt solution $\hat{\rho}_*$ concentrates on a ball of radius one around the origin,

$$\frac{2m}{m-1}\hat{\rho}_*(\hat{x})^{m-1} \;=\; \left(1 - |\hat{x}|^2\right)_+. \tag{4}$$

This normalization will simplify the notation in the subsequent sections. In the sequel, we will often refer to $\hat{\rho}_*$ as the *Barenblatt profile*.

The advantage of working with (3) instead of (1) relies on the fact that the above rescaling comes along with a change of perspective: the Barenblatt solution now becomes a fixed point of the equation. In fact, as we will see below, for fixed total mass $\hat{M}$, it is both the unique stationary solution of (3) and the ground state of the physical entropy

$$\hat{E}(\hat{\rho}) \;=\; \frac{1}{m-1}\int \hat{\rho}(\hat{x})^m \, d\hat{x} + \frac{1}{2}\int |\hat{x}|^2 \hat{\rho}(\hat{x}) \, d\hat{x}. \tag{5}$$

In this regard, the porous medium equation (1) shows a substantially different behavior than the confined porous medium equation (3): There is no stationary solution of (1) under the mass constraint $\|\rho\|_{L^1} = M$ for any $M > 0$. In other words, solutions to the porous medium equation do not relax in finite time and the ground state can be attained only asymptotically in the long-time limit.

Somehow related to this feature is the fact that, opposed to the original equation (1), the confined porous medium equation (3) is no longer invariant under spatial translations. More precisely, in the new variables the origin is the unique attraction point in $\mathbb{R}^N$ around which the mass asymptotically concentrates, driven by the convection term $-\hat{\nabla}\cdot(\hat{x}\hat{\rho})$. Due to this loss of translation invariance it is therefore not surprising that those initial data $\hat{\rho}_0$ that differ from the ground state $\hat{\rho}_*$ only by a shift, $\hat{\rho}_0(\hat{x}) = \hat{\rho}_*(\hat{x} - \hat{x}_0)$ for some $\hat{x}_0 \in \mathbb{R}^N$, will play a distinguished role in the discussion of our result on refined asymptotics of (3): Spatial translations correspond to the smallest eigenvalue of the linearized operator and generate thus those solutions that saturate the optimal rate of convergence.

The above rescaling, however, also has a small drawback: It is singular at time zero. To compare "initial data", one would better consider the moment $t = 1$. More precisely, up to spatial dilations both in the x and ρ variables, the initial datum $\hat{\rho}(\hat{t} = 0, \cdot\,)$ corresponds to the original solution at time one $\rho(t = 1, \cdot\,)$. Because of our interest in the long-time dynamics, this skew correspondence, however, has no influence neither on the subsequent analysis nor on the interpretation of our result.

It remains to discuss the stationarity of $\hat{\rho}_*$ for the nonlinear Fokker–Planck equation (3) and its minimality for the entropy (5). Calculating the energy dissipation along trajectories of the flow (3),

$$\frac{d}{d\hat{t}}\hat{E}(\hat{\rho}(\hat{t})) \;=\; -\int \hat{\rho}\left|\hat{\nabla}\left(\frac{1}{2}|\hat{x}|^2 + \frac{m}{m-1}\hat{\rho}^{m-1}\right)\right|^2 d\hat{x},$$

we see that the dissipation rate is zero if and only if $\frac{1}{2}|\hat{x}|^2 + \frac{m}{m-1}\hat{\rho}^{m-1}$ is constant, that is, if and only if $\hat{\rho}$ is of the form (4), provided that $\hat{\rho}$ and $\hat{\rho}_*$ have the same total mass. This implies that $\hat{\rho}_*$ is the unique stationary solution of (3). To deduce the minimality of $\hat{\rho}_*$ for the entropy $\hat{E}$, we additionally observe that the entropy is strictly convex on the convex configuration space of nonnegative densities with fixed total mass $\hat{M}$, and admits thus a unique minimizer.

1.2 Otto's gradient flow interpretation and the Wasserstein distance

In his seminal paper [22], Otto introduced a new and physically meaningful gradient flow interpretation of the porous medium equation (1). This interpretation is based on the Lyapunov approach of Newman & Ralston in [20, 27]. In the abstract setting, a gradient flow is the dynamical system on a Riemannian manifold $(\mathcal{M}, g)$ that describes the evolution as the steepest descent in an energy landscape, i.e.,

$$\partial_t \rho + \operatorname{grad} E(\rho) \;=\; 0. \tag{6}$$

Here, the gradient $\operatorname{grad} E$ of the energy E is defined via Riesz' representation theorem

$$\operatorname{diff} E(\rho).\delta\rho \;=\; g_\rho(\operatorname{grad} E(\rho), \delta\rho) \quad \text{for all } \rho \in T_\rho\mathcal{M},$$

where $\operatorname{diff} E$ denotes the differential of E and $T_\rho\mathcal{M}$ is the tangent plane at the point $\rho \in \mathcal{M}$. A short formal computation shows that (6) can be equivalently stated as

$$\partial_t \rho \;=\; \operatorname{argmin}\left\{ \frac{1}{2} g(\delta\rho, \delta\rho) + \operatorname{diff} E(\rho).\delta\rho \;:\; \delta\rho \in T_\rho\mathcal{M} \right\}.$$

In the following, we identify the gradient flow ingredients $\mathcal{M}$, g, and E that constitute the porous medium equation using Otto's approach. (In fact, different gradient flow interpretations were proposed for the porous medium equation. Otto's gradient flow is natural in view of the thermodynamical background of the equation. Moreover, it also pertains to the confined evolution, whereas the traditional approaches do not.)

As solutions to (1) preserve non-negativity (in fact, solutions satisfy a comparison principle) and the total mass M, the manifold in the gradient flow interpretation of the porous medium equation has to be chosen as

$$\mathcal{M} \;=\; \left\{ \rho \;:\; \rho \geq 0 \text{ and } \int \rho \, dx = M \right\}.$$

Tangent fields $\delta\rho$ at $\rho \in \mathcal{M}$ must be mass-preserving and non-negative where $\rho = 0$. However, as the evolution (1) does not allow for nucleation of mass outside of the support of ρ, i.e., $\operatorname{supp}(\partial_t \rho) \subset \operatorname{supp}(\rho)$, it is natural to think of the tangent plane as

$$T_\rho\mathcal{M} \;=\; \left\{ \delta\rho \;:\; \operatorname{supp}(\delta\rho) \subset \operatorname{supp}(\rho) \text{ and } \int \delta\rho = 0 \right\}.$$

Observe that this restriction of tangent fields does not suppress the motion of the free boundary since $\delta\rho$ may be nonzero on $\partial \operatorname{supp}(\rho)$. The metric tensor is defined by

$$g_\rho(\delta\rho, \delta\rho) \;=\; \int \rho |\nabla\psi|^2 \, dx,$$

where ψ and $\delta\rho$ are related via the elliptic boundary value problem

$$\begin{aligned} -\nabla\cdot(\rho\nabla\psi) &= \delta\rho \qquad \text{in } \operatorname{supp}(\rho), \\ \rho\nabla\psi\cdot\nu &= 0 \qquad \text{on } \partial\operatorname{supp}(\rho). \end{aligned} \tag{7}$$

Here, ν denotes the outer normal vector on $\partial\operatorname{supp}\rho$. Actually, arguing even more formally, Otto set the elliptic equation in the entire space $\mathbb{R}^N$, cf. [22, eq. (8)]. However, in order to motivate our later analysis, here, we equip the equation with its natural boundary conditions. That is, assuming enough regularity of ρ and $\delta\rho$, the distributional solution ψ of $-\nabla\cdot(\rho\nabla\psi)=\delta\rho$ satisfies (7). Again, we remark that $\nabla\psi$ may be non-zero on $\partial\operatorname{supp}(\rho)$, so that the boundary is indeed "free" fo move. Equivalently, we can characterize the metric tensor variationally:

$$\frac{1}{2}g_\rho(\delta\rho,\delta\rho) \;=\; \sup_\varphi\left\{-\int\rho|\nabla\varphi|^2\,dx-\int\varphi\delta\rho\,dx\right\},$$

cf. [23, eq. (2.9)], where the supremum is taken over all smooth functions φ on $\mathbb{R}^N$. The optimal φ is this formulation satisfies (7). By identifying tangent fields $\delta\rho$ with these new variables, the tangent plane at ρ can be rewritten as the weighted Sobolev space

$$T_\rho\mathcal{M} \;=\; \left\{\psi\;:\;\int\rho|\nabla\psi|^2\,dx<\infty\right\}. \tag{8}$$

In particular, the metric tensor is finite on tangent fields.

The energy functional in the gradient flow interpretation is given by the entropy

$$E(\rho) \;=\; \frac{1}{m-1}\int\rho^m\,dx. \tag{9}$$

A short computation now shows the porous medium equation (1) is indeed the gradient flow for $(\mathcal{M},g)$ and E defined as above, see [22, Sec. 1.3] for details.

As for the confined porous medium equation (3), we remark that $\hat\rho$ still evolves according to the gradient flow on the same Riemannian manifold as ρ, now with total mass $\hat M$ and denoted by $(\hat{\mathcal{M}},\hat g)$, but for the energy $\hat E$ defined in (5).

We finally address the induced geodesic distance on the Riemannian manifold $(\hat{\mathcal{M}},\hat g)$. Benamou & Brenier [5] discovered the relation between the Kantorovich mass transfer problem and continuum mechanics by identifying the geodesic distance on $(\hat{\mathcal{M}},\hat g)$ between two configurations $\hat\rho_0$, $\hat\rho_1$ having finite second moments $\int|\hat x|^2\hat\rho_i\,d\hat x<\infty$ with the *Wasserstein distance*

$$d_2(\hat\rho_1,\hat\rho_0)^2 \;=\; \inf_{\mu\in\Gamma(\hat\rho_1,\hat\rho_0)}\iint|\hat x_1-\hat x_0|^2\,\mu(d\hat x_1 d\hat x_0),$$

see also [22, Sec. 4.3] or [23, Prop. 4.3]. Here the infimum is taken over the space $\Gamma(\hat\rho_1,\hat\rho_0)$ of all nonnegative measures μ on $\mathbb{R}^N\times\mathbb{R}^N$ having marginals $\hat\rho_1$ and $\hat\rho_0$, i.e.,

$$\iint\zeta(\hat x_i)\mu(d\hat x_1 d\hat x_0)=\int\zeta(\hat x_i)\;\hat\rho_i(\hat x_i)d\hat x_i,$$

for all $\zeta \in C_c(\mathbb{R}^d)$ and $i = 0, 1$. For a detailed study of Wasserstein distances and more general optimal transportation distances, we refer the interested reader to Villani's two monographs [32, 33].

1.3 Linearization

Instead of linearizing the confined porous medium equation (3) around the Barenblatt profile $\hat\rho_*$, we imitate the approach of Denzler & McCann [9] and compute the Hessian of the entropy $\hat E$ at $\hat\rho_*$. In view of the gradient flow formulation described in the previous subsection, this is formally equivalent to linearizing (3) directly, but it has the advantage that a natural class of perturbations is intrinsically given by the tangent fields.

Tangent fields $\delta\hat\rho$ are mass-preserving and concentrate on the support of $\hat\rho_*$. Following Otto, and as described in the previous subsection, we identify such tangent fields with new variables ψ via the elliptic problem (7), i.e.,

$$\begin{aligned} -\nabla\cdot(\hat\rho_*\nabla\psi) &= \delta\hat\rho \quad \text{in } B_1, \\ \hat\rho_*\hat\nabla\psi\cdot\nu &= 0 \quad \text{on } \partial B_1, \end{aligned} \tag{10}$$

where B_1 denotes the ball of radius one centered at the origin, and thus $B_1 = \operatorname{supp}(\hat\rho_*)$. Moreover, ψ is such that

$$\int \hat\rho_*|\hat\nabla\psi|^2\,d\hat x < \infty, \tag{11}$$

cf. (8). Geodesic curves $\{\hat\rho_s\}_{|s|\ll 1}$ on $\mathcal{M}$ passing through $\hat\rho_*$ and pointing in the direction $\delta\hat\rho$ are obtained by McCann's [19] displacement interpolation

$$\hat\rho_*(\hat x) \;=\; \det\Big(I + s\hat\nabla^2\psi(\hat x)\Big)\,\hat\rho_s\Big(\hat x + s\hat\nabla\psi(\hat x)\Big) \tag{12}$$

(see also [22, Sec. 4.3]), that is, $\hat\rho_*$ is pushed forward by the flux $\mathrm{id} + s\hat\nabla\psi(\,\cdot\,)$. We note that

$$\delta\hat\rho \;=\; \partial_s|_{s=0}\,\hat\rho_s \;=\; -\hat\nabla\cdot\Big(\hat\rho_*\hat\nabla\psi\Big).$$

Recalling that the Hessian $\operatorname{hess}\hat E$ of a function $\hat E$ on a Riemannian manifold $(\hat{\mathcal{M}},\hat g)$ can be calculated as the second derivative of $\hat E$ along geodesics, we define and have

$$\operatorname{Hess}\hat E(\hat\rho_*)(\psi,\psi) \;:=\; \hat g_{\hat\rho_*}(\delta\hat\rho, \operatorname{hess}\hat E(\hat\rho_*)\delta\hat\rho) \;=\; \frac{d^2}{ds^2}\bigg|_{s=0}\hat E(\hat\rho_s).$$

A *formal* computation yields:

$$\begin{aligned} &\operatorname{Hess}\hat E(\hat\rho_*)(\psi,\psi) \\ &\quad= \; m\int \hat\rho_*^{m-2}\,(\partial_s|_{s=0}\,\hat\rho_s)^2\;d\hat x + \int\left(\frac{m}{m-1}\hat\rho_*^{m-1} + \frac12|\hat x|^2\right)\partial_s^2|_{s=0}\,\hat\rho_s\,d\hat x. \end{aligned}$$

Observe that $\operatorname{supp} \partial_s^2|_{s=0} \hat{\rho}_s \subset \operatorname{supp} \hat{\rho}_*$ and $\int \partial_s^2|_{s=0} \hat{\rho}_s \, d\hat{x} = 0$, and so the second integral on the right vanishes thanks to the definition of the Barenblatt profile $\hat{\rho}_*$. Because of $\partial_s|_{s=0} \hat{\rho}_s = -\hat{\nabla} \cdot \left(\hat{\rho}_* \hat{\nabla} \psi \right)$, we can rewrite the Hessian as

$$\operatorname{Hess} \hat{E}(\hat{\rho}_*)(\psi, \psi) \;=\; m \int \hat{\rho}_*^{m-2} \left(\hat{\nabla} \cdot \left(\hat{\rho}_* \hat{\nabla} \psi \right) \right)^2 d\hat{x}.$$

Finally, integration by parts yields

$$\operatorname{Hess} \hat{E}(\hat{\rho}_*)(\psi, \psi) \;=\; \int \hat{\rho}_* \hat{\nabla} \psi \cdot \hat{\nabla} \left(-m \hat{\rho}_*^{m-2} \hat{\nabla} \cdot \left(\hat{\rho}_* \hat{\nabla} \psi \right) \right) d\hat{x}.$$

At this point, we consider the derivation of the Hessian on a purely formal level. The above computations certainly hold true for functions ψ that are smooth up to the boundary. In this case, the integrals are well defined and we can integrate by parts since $\hat{\rho}_*^{m-1}$ vanishes on $\partial \operatorname{supp}(\hat{\rho}_*)$.

We finally complete this subsection by introducing the *displacement Hessian*

$$\mathcal{H} \psi \;=\; -m \hat{\rho}_*^{m-2} \hat{\nabla} \cdot \left(\hat{\rho}_* \hat{\nabla} \psi \right). \tag{13}$$

In view of the explicit formula (4) for $\hat{\rho}_*$, the displacement Hessian can be rewritten as

$$\mathcal{H} \psi(x) \;=\; -\frac{m-1}{2} \left(1 - |\hat{x}|^2 \right) \hat{\Delta} \psi(\hat{x}) + \hat{x} \cdot \hat{\nabla} \psi(\hat{x}) \quad \text{for } \hat{x} \in B_1. \tag{14}$$

With the help of $\mathcal{H}$, the linearized confined porous medium equation reads $\partial_{\hat{t}} \psi + \mathcal{H} \psi = 0$, or on the level of $\delta \rho$:

$$\partial_{\hat{t}} \delta \hat{\rho} + \mathcal{L}^{-1} \mathcal{H} \mathcal{L} \delta \hat{\rho} \;=\; 0, \tag{15}$$

where the operator $\mathcal{L}$ is defined via $\mathcal{L} \delta \hat{\rho} = \psi$ and $\delta \hat{\rho}$ and ψ are related in the usual way. Moreover, a formal analysis yields that $\mathcal{H}$ and hess are similar in the sense that $\text{hess} = \mathcal{L}^{-1} \mathcal{H} \mathcal{L}$.

The remainder of the paper is exclusively devoted to the analysis of the displacement Hessian. We will show in Proposition 1 below, that it defines a self-adjoint operator on a suitable domain $\mathcal{D}(\mathcal{H})$. Our main result, stated in Theorem 1 below, gives the complete spectrum of $\mathcal{H}$.

2 Rigorous part

From now on we claim mathematical rigor. As we focus exclusively on the confined porous medium equation (3) we may simplify our notation and drop the hats from the rescaled variables. Moreover, we restrict to the case

$$N \geq 2.$$

Let $H^1_{\rho_*}$ denote the class of all locally integrable functions on $B_1 = \operatorname{supp}\rho_*$ such that (11) holds, i.e.,

$$\|\psi\|^2_{H^1_{\rho_*}} := \int \rho_* |\nabla\psi|^2 \, dx < \infty,$$

with the identification of two functions if they only differ by a constant. Observe that this space is a separable Hilbert space for the topology induced by the norm $\|\cdot\|_{H^1_{\rho_*}}$. In the appendix, we study some properties related to elliptic theory in $H^1_{\rho_*}$.

So far, the derivation of the Hessian of the energy functional at the Barenblatt profile and the definition of the displacement Hessian were only formally justified. Hence, as a first step before debarking into spectral analysis, we have to build a rigorous basis for our investigations. Since the formal computations of the preceding section are certainly valid for functions that are smooth up to the boundary of the support of the Barenblatt profile, i.e., $C^\infty(\bar{B}_1)$ functions, let us denote the displacement Hessian defined in (13) by $\mathcal{H}|_{C^\infty(\bar{B}_1)}$. A short computation yields that this operator is nonnegative and symmetric (cf. proof of Proposition 1). In Subsection 2.2 below, we show that $\mathcal{H}|_{C^\infty(\bar{B}_1)}$ extends naturally to a nonnegative self-adjoint operator $\mathcal{H}$ with domain

$$\begin{aligned}\mathcal{D}(\mathcal{H}) = \big\{ &\psi \in H^3_{loc}(B_1) \cap H^1_{\rho_*} : \\ &\rho_*^{m-1}\nabla\cdot(\rho_*\nabla\psi) \in H^1_{\rho_*},\ \rho_*\nabla\psi\cdot\nu = 0 \text{ on } \partial B_1 \big\}. \end{aligned} \tag{16}$$

Here, the condition $\rho_*\nabla\psi\cdot\nu = 0$ on ∂B_1 means that

$$\int \rho_*\nabla\xi\cdot\nabla\psi\,dx \;=\; -\int \xi\nabla\cdot(\rho_*\nabla\psi)\,dx \quad \text{for all } \xi \in H^1_{\rho_*}. \tag{17}$$

We also remark that the regularity that is assumed in the definition of $\mathcal{D}(\mathcal{H})$ implies that $\mathcal{H}\psi = -m\rho_*^{m-1}\nabla\cdot(\rho_*\nabla\psi)$ for all $\psi \in \mathcal{D}(\mathcal{H})$, where the derivatives have to be understood in the weak Sobolev sense.

Remark 1. *The boundary conditions in* (17) *are* asymptotic *boundary conditions in the sense that*

$$\lim_{|x|\uparrow 1} \rho_*(x)\nabla\psi(x)\cdot\frac{x}{|x|} = 0, \tag{18}$$

for every function $\psi \in C^\infty(\bar{B}_1)$. In fact, thanks to the density of $C^\infty(\bar{B}_1)$ functions in $H^1_{\rho_*}$ by Lemma 2 in the appendix, and the Sobolev embedding $H^1_{\rho_*} \subset L^2(B_1, \rho_*^{2-m}dx)$ in Lemma 3 in the appendix, it is enough to restrict condition (17) *to functions $\xi \in C^\infty(\bar{B}_1)$. Then, the equivalence of* (17) *and* (18) *follows immediately (for instance, by an indirect argument).*

2.1 Main results

We are now on the position to state our main results:

Theorem 1. *The operator $\mathcal{H} : \mathcal{D}(\mathcal{H}) \to H^1_{\rho_*}$ is self-adjoint. Its spectrum consists only of eigenvalues, given by*

$$\lambda_{\ell k} \;=\; \ell + 2k + 2k(k + \ell + \frac{N}{2} - 1)(m-1),$$

for ℓ, $k \in \mathbb{N} \cup \{0\}$ with $(\ell, k) \neq (0,0)$. The corresponding eigenfunctions form an orthogonal basis of $H^1_{\rho_*}$ and are given by polynomials of the form

$$\psi_{\ell m k}(x) = F(-k, \frac{1}{m-1} + \ell + \frac{N}{2} - 1 + k; \ell + \frac{N}{2}; |x|^2) Y_{\ell m}\left(\frac{x}{|x|}\right) |x|^\ell,$$

$m \in \{1, \dots, M_\ell\}$, where $F(a, b; c; z)$ is a hypergeometric function and $Y_{\ell m}$ is a spherical harmonic, corresponding to the eigenvalue $\ell(\ell + N - 2)$ of $-\Delta_{\mathbb{S}^{N-1}}$ with multiplicity $M_\ell = \frac{(N+\ell-3)!(N+2\ell-2)}{\ell!(N-2)!}$.

The eigenfunctions corresponding to the eigenvalues $\lambda_{\ell k}$ are polynomials of degree $\ell + 2k$, which are harmonic and homogeneous if $k = 0$. In fact, the hypergeometric functions $F(a, b; c; z)$ in the statement of the theorem reduce to polynomials of degree k in the case $a = -k$, see Subsection 2.4 below. Spherical harmonics $Y_{\ell m}$ are harmonic homogeneous polynomials of degree ℓ. These functions are ubiquitous in mathematical physics, see e.g. [13] to name but a single reference.

We observe that the eigenvalues of the displacement Hessian $\mathcal{H}$ are affine functions of the parameter m. There are countably many constant levels $\lambda_{\ell 0} = \ell$. Above λ_{10} and λ_{20}, the spectrum features a crossing of eigenvalues when varying m. A first level crossing occurs for the eigenvalues $\lambda_{30} = 3$ and $\lambda_{01} = 2 + N(m-1)$ at $N(m-1) = 1$, and the number of crossings increases with ℓ. The eigenvalues coincide with the eigenvalues found by Denzler & McCann [9] for the fast diffusion regime in the sense that each eigenvalue continues to the range $m < 1$ if m is sufficiently close to 1. For smaller m, the eigenvalues dissolve into continuous spectrum (that disappears in the limit $m \uparrow 1$). The occurrence of a continuous spectrum for the fast diffusion equation is analytically related to the fact that in the fast-diffusion regime, the number of moments possessed by the Barenblatt profile is finite — in contrast to the situation for the porous medium equation. In the limit $m \downarrow 1$, we recover the spectrum of the Ornstein–Uhlenbeck operator $-\Delta + x \cdot \nabla$. This coincidence is not surprising since the Ornstein–Uhlenbeck operator is the limit operator of $\mathcal{H}$ as $m \downarrow 1$. In this case, ρ_* is a Gaussian.

The knowledge of the spectrum of the displacement Hessian $\mathcal{H}$ suggests the asymptotic expansion to leading order of solutions $\rho = \rho(t)$ to the confined porous medium equation close to the stationary solution ρ_*. Arguing purely formally again, we recall the operator $\mathcal{L}$ defined by $\mathcal{L}\delta\rho = \psi$ where $\delta\rho$ and ψ are related in the usual way (10), cf. Subsection 1.3, and observe that the spectrum of $\mathcal{H}$ coincides with the spectrum of $\mathcal{L}^{-1}\mathcal{H}\mathcal{L}$. Moreover, the corresponding eigenfunctions $\delta\rho_{\ell k m} = \mathcal{L}^{-1}\psi_{\ell k m}$ form an orthogonal basis of the Hilbert space $L^2(B_1, \rho_*^{m-2} dx)$. For solutions $\rho(t)$ close to ρ_*, equation (3) reads to leading order

$$\partial_t(\rho(t) - \rho_*) \;\approx\; -\mathcal{L}^{-1}\mathcal{H}\mathcal{L}(\rho(t) - \rho_*),$$



Figure 1: 2D plots of the supports of the affine transformations (dashed and dotted lines) of the Barenblatt profile (solid lines) generated by the eigenfunctions corresponding to the eigenvalues λ_{10}, λ_{20}, λ_{30}, and λ_{01} (from the left to the right).

cf. (15), and thus, exploiting the knowledge of the complete spectrum of $\mathcal{H}$ (and thus of $\mathcal{L}^{-1}\mathcal{H}\mathcal{L}$)

$$\begin{aligned}\rho(t) - \rho_* \\ = \sum_{\ell k m} c_{\ell k m} \delta\rho_{\ell k m} e^{-\lambda_{\ell k} t} + \sum_{\ell k m} \sum_{\ell' k' m'} c_{\ell k m, \ell' k' m'} \delta\rho_{\ell k m} \delta\rho_{\ell' k' m'} e^{-(\lambda_{\ell k} + \lambda_{\ell' k'})t} + \dots,\end{aligned}$$

where the c's are constants that depend on the initial data $\rho(0)$ only. In order to investigate the precise asymptotics of the nonlinear equation (3), one has to take into account terms of higher order in $\rho(t) - \rho_*$. This problem will be addressed in future research. In this context we should also mention the work of Koch [18, Ch. 5.3.4], which offers a framework which seems to be suitable for a rigorous treatment of higher order asymptotics. For the one-dimensional porous medium equation, the full asymptotics was rigorously computed by Angenent in [2], based on an earlier investigation by Zel'dovic & Barenblatt [34]. This present work may be considered as a first attempt to complement Angenent's result in the multivariable case. Recently, Denzler, Koch & McCann studied the higher order asymptotic behavior of solutions to the fast-diffusion equation [8], using subtle dynamical systems arguments in weighted Hölder spaces.

The main purpose of understanding the full asymptotic expansion of solutions in terms of the eigenfunctions of the Hessian around the self-similar solution is to get a deeper understanding of the long-time behavior of solutions. Although we have to keep in mind that our investigation is not fully rigorous in the derivation of the displacement Hessian operator, we will in the following discuss the the role of the smallest eigenvalues and their corresponding eigenfunctions for the convergence towards the attractor ρ_*. We recall that the geodesic curves in the Wasserstein space passing through the Barenblatt solution ρ_* along the vector field $\nabla\psi$ are characterized by $\rho_*(x) = \det(I + s\nabla^2\psi)\,\rho_s(x + s\nabla\psi)$ (cf. (12)). With this formula at hand, the interpretation of the eigenfunctions as generators of mass transport is immediate. The smallest eigenvalue $\lambda_{10} = 1$ corresponds to transitions along the axis e_m with $m \in \{1, \dots, M_1 = N\}$ (see Figure 1a). Indeed, in this case, the eigenfunction is a homogeneous polynomial of degree one, and the displacement is

generated by affine perturbations of the identity map $x + sc_m e_m$ (here, c_m denotes a normalizing factor). This fact is in good agreement with the optimal convergence rate of solutions of the porous medium equation to the Barenblatt profile,

$$d_2(\rho(t), \rho_*) \leq e^{-t} d_2(\rho_0, \rho_*),$$

here stated in the Wasserstein topology [22, Theorem 1]. In fact, spatially translated Barenblatt profiles saturate this bound. The eigenfunctions to the second smallest eigenvalue $\lambda_{20} = 2$ generate affine transformations on $\mathbb{R}^N$ modulo rotations (Figure 1b). To see this, we observe that ψ_{20m} is a harmonic homogeneous polynomial of degree two, and thus $x + s\nabla\psi = x + sA_{2m}x$ for some symmetric and trace-free matrix A_{2m}. In other words, solutions to (3) generated by affine transformations show a rate of convergence of precise order e^{-2t} to the Barenblatt profile. Denzler & McCann [10] give an exact description of the invariant manifold corresponding to these affine perturbations. The role of the next term in the expansion depends on the value of $(m-1)N$. As indicated above, there is a first level crossing of the eigenvalues $\lambda_{\ell 0}$ and λ_{01}: It is $\lambda_{01} \leq \lambda_{\ell 0}$ precisely for $\ell \geq 1/\alpha$. Eigenfunctions to the eigenvalue λ_{01} corresponds to isotropic dilations of the Barenblatt profile (see Figure 1d). Indeed, the spherical harmonic Y_{01} degenerates to a constant and ψ_{011} is proportional to $1 - (\alpha(m-1)N)^{-1}|x|^2$. It follows that $x + s\nabla\psi = (1 - cs)x$ for some constant c. Such solutions converge with a rate $e^{-t/\alpha}$. The mass transport corresponding to the $(\ell, 0)$-modes with $\ell \geq 3$ is less obvious, as the corresponding eigenfunctions $\psi_{\ell 0m}$ are homogenous harmonic polynomials of degree ℓ. We only discuss the situation where $\ell = 3$. In this case, the transformation is quadratic and leads to triangular deformations (see Figure 1c). The rate of convergence is e^{-3t}. The deformations corresponding to the remaining eigenvalues can be similarly computed although the complexity of the underlying symmetries is increasing.

We finally remark that the knowledge of spectral properties of the displacement Hessian $\mathcal{H}$, and of the smallest eigenvalue $\lambda_{10} = 1$ in particular, immediately yields the sharp spectral gap estimate

$$\int \rho_* |\nabla\psi|^2 \, dx \;\leq\; \operatorname{Hess} E(\rho_*)(\psi, \psi),$$

for all $\psi \in H^1_{\rho_*}$. This estimate, was already derived by Otto [22, Sec. 4.4] and builds on McCann's displacement convexity [19].

We prove Theorem 1 in the remainder of this article: In subsection 2.2 we show that the displacement Hessian $\mathcal{H}$ can be understood as a self-adjoint operator on $H^1_{\rho_*}$ (Proposition 1). Moreover, we show that its spectrum is purely discrete provided that $\mathcal{H}$ has infinitely many eigenvalues that accumulate at infinity (Proposition 2). The proof of the latter fact requires some preparation. In Subsection 2.3 we make a change of variables into spherical coordinates, which has the advantage that the eigenvalue problem reduces to a one-dimensional problem for the radial components of the displacement Hessian. This eigenvalue problem is solved in Subsection 2.4. We

find that the radial components of $\mathcal{H}$ have indeed countably many eigenvalues (and so has $\mathcal{H}$). These eigenvalues and the corresponding eigenfunctions are explicitly computed in Proposition 3.

2.2 Self-adjointness and spectral decomposition of $\mathcal{H}$

Since $\mathcal{H}|_{C^\infty(\bar{B}_1)}$ is nonnegative and symmetric (cf. proof of Proposition 1) and because $C^\infty(\bar{B}_1)$ is densely contained in $H^1_{\rho_*}$ by Lemma 2 in the appendix, the operator $\mathcal{H}|_{C^\infty(\bar{B}_1)}$ is closable in $H^1_{\rho_*}$ and its closure is still nonnegative and symmetric. We denote this closure again by $\mathcal{H}$. In the following we show that $\mathcal{H}$ is self-adjoint with domain $\mathcal{D}(\mathcal{H})$ defined in (16).

The following proposition contains a first statement on the spectrum of $\mathcal{H}$, namely that the spectrum is real and contained in the ray $(0, \infty)$. The first fact is a consequence of self-adjointness, the second one of nonnegativity together with the fact that $\mathcal{H}$ is one-to-one and onto (and thus 0 must be in the resolvent set).

Proposition 1 (Self-adjointness)**.** *The operator $\mathcal{H} : \mathcal{D}(\mathcal{H}) \to H^1_{\rho_*}$ is nonnegative, one-to-one and onto. In particular, $\mathcal{H}$ is self-adjoint.*

In fact, it can be shown that $\mathcal{H}$ is the only self-adjoint extension of $\mathcal{H}|_{C^\infty(\bar{B}_1)}$, cf. [9, Lemma 4]. For the theory of unbounded operators in general and self-adjointness in particular we refer to [28, Ch. 13].

The following proposition anticipates some of the results in Proposition 3 below, namely that $\mathcal{H}$ has countably many eigenvalues that accumulate at infinity. Supposing this, we combine some properties of $\mathcal{H}$ with general results from functional analysis to deduce that the spectrum of $\mathcal{H}$ is purely discrete. As a consequence, it is enough to solve the eigenvalue problem in order to compute the complete spectrum. This will be our concern in the two subsections that follow.

Proposition 2 (Discrete spectrum)**.** *Suppose that the displacement Hessian $\mathcal{H} : \mathcal{D}(\mathcal{H}) \to H^1_{\rho_*}$ has countably many eigenvalues that accumulate at infinity. Then the eigenfunctions of $\mathcal{H}$ form an orthogonal basis of $H^1_{\rho_*}$. In particular, the spectrum of $\mathcal{H}$ is purely discrete, i.e., it consists only of eigenvalues with finite multiplicity and which do not have a finite accumulation point.*

We start with the

Proof of Proposition 1. Since $C^\infty(\bar{B}_1)$ is dense in $H^1_{\rho_*}$ by Lemma 2 and $C^\infty(\bar{B}_1) \subset \mathcal{D}(\mathcal{H}) \subset H^1_{\rho_*}$, we immediately have that $\mathcal{H}$ is densely defined. We first show that $\mathcal{H}$ is a nonnegative, symmetric operator on $\mathcal{D}(\mathcal{H})$. Indeed, by the definition of $\mathcal{H}$ and by integration by parts we have

$$\int \rho_* \nabla\varphi \cdot \nabla\left(\mathcal{H}\psi\right) dx \;=\; m \int \rho_*^{m-2} \nabla \cdot (\rho_* \nabla\varphi) \nabla \cdot (\rho_* \nabla\psi)\, dx$$

for all $\varphi, \psi \in \mathcal{D}(\mathcal{H})$. Notice that the boundary term vanishes when integrating by parts because of (17) and $\mathcal{H}\psi \in H^1_{\rho_*}$. Now nonnegativity follows from choosing

$\varphi = \psi$ and symmetry comes from the symmetry in φ and ψ of the right hand side of the above identity.

Notice also that a densely defined, symmetric operator is automatically one-to-one and self-adjoint if it is onto, cf. [28, Theorem 13.11]. In our situation, the latter is a consequence of Lemmas 3 & 5 in the appendix. Indeed, given $u \in H^1_{\rho_*}$, we may without loss of generality assume that $\int \rho_*^{2-m} u \, dx = 0$ as we identify functions that are the same up to a constant. By Lemma 3 we have that $u \in L^2(B_1, \rho_*^{2-m} dx)$. We define $\tilde{u} = \frac{1}{m}\rho_*^{2-m} u$ and observe that this function satisfies the hypotheses of Lemma 5. Consequently, there exists a function $\psi \in H^2_{\mathrm{loc}} \cap H^1_{\rho_*}$ such that

$$\begin{aligned} -\nabla \cdot (\rho_* \nabla \psi) &= \tilde{u} \quad \text{in } B_1, \\ \rho_* \nabla \psi \cdot \nu &= 0 \quad \text{on } \partial B_1. \end{aligned}$$

By the definition of $\tilde{u}$, the equation in the ball can be rewritten as $-m\rho_*^{m-2}\nabla \cdot (\rho_* \nabla \psi) = u$, i.e., $\mathcal{H}\psi = u$. Higher interior regularity estimates (cf. [28, Theorem 8.12]) moreover show that $\psi \in H^3_{\mathrm{loc}} \cap H^1_{\rho_*}$. Hence $\psi \in \mathcal{D}(\mathcal{H})$. This proves that $\mathcal{H}$ is onto, which completes the proof of Proposition 1. □

We finally prove Proposition 2.

Proof of Proposition 2. We first show that the first statement implies the second one, and thus we assume that the eigenfunctions of $\mathcal{H}$ form an orthogonal basis of $H^1_{\rho_*}$. Then the operator $\mathcal{H}$ can be written as $\mathcal{H} = \sum_n \lambda_n \mathcal{P}_n$, where the λ_n's are the eigenvalues of $\mathcal{H}$ and the $\mathcal{P}_n$'s the orthogonal projections onto the corresponding eigenspaces. We define $dE = \sum_n \mathcal{P}_n \delta_n$ with δ_n being the delta measure concentrated at the point λ_n, that is, for every Borel set $A \subset \mathbb{R}$, we have

$$E(A) = \int_A dE(\lambda) = \sum_{\lambda_n \in A} \mathcal{P}_n.$$

One easily checks that E defines a spectral measure. Moreover, since

$$\mathcal{H} = \int_{-\infty}^{+\infty} \lambda \, dE(\lambda),$$

the spectral measure E is the unique spectral decomposition of $\mathcal{H}$ (cf. [28, Theorem 13.30]). Now the discreteness of E implies the discreteness of the spectrum of $\mathcal{H}$.

To prove that $H^1_{\rho_*}$ admits indeed an orthogonal basis of eigenfunctions of $\mathcal{H}$, it is enough to show that $\mathcal{H}$ has a compact self-adjoint inverse $\mathcal{H}^{-1} : H^1_{\rho_*} \to H^1_{\rho_*}$. In fact, in view of the spectral theorem for compact operators on separable Hilbert spaces, cf. [28, Theorem 4.25], the spectrum of $\mathcal{H}^{-1}$ is at most countable. Since the eigenvalue problems for $\mathcal{H}^{-1}$ and $\mathcal{H}$ are equivalent with reciprocal eigenvalues and $\mathcal{H}$ has countably many eigenvalues with accumulation point at infinity by assumption, the spectrum of $\mathcal{H}^{-1}$ must also be countable, and thus, the eigenfunctions form an orthogonal basis, cf. [28, Theorem 12.29].

We now show that $\mathcal{H}$ has a compact self-adjoint inverse. As $\mathcal{H}$ is one-to-one and onto, its inverse $\mathcal{H}^{-1} : H^1_{\rho_*} \to H^1_{\rho_*}$ is well-defined, that is, given $\xi \in H^1_{\rho_*}$, we define $\mathcal{H}^{-1}\xi = \psi$, where ψ is the solution of

$$\begin{aligned} -m\rho_*^{m-2}\nabla\cdot(\rho_*\nabla\psi) &= \xi \quad \text{in } B_1, \\ \rho_*\nabla\psi\cdot\nu &= 0 \quad \text{on } \partial B_1. \end{aligned}$$

In the definition of the Hilbert space $H^1_{\rho_*}$, we identified functions that only differ by a constant. In the following, we fix constants by additionally requiring that

$$\int \rho_*^{2-m}\psi\,dx \;=\; \int \rho_*^{2-m}\xi\,dx \;=\; 0.$$

Notice also that $\mathcal{H}^{-1}$ is bounded and self-adjoint because $\mathcal{H}$ is self-adjoint, one-to-one and onto by Proposition 1, cf. [28, Theorem 13.11].

It remains to show that $\mathcal{H}^{-1}$ is a compact operator. The argument is standard, and is based on a Rellich lemma, namely Lemma 4 in the appendix. We display the argument since we are dealing with weighted (and thus non-standard) Sobolev norms. Let $\{\xi_n\}_{n\in\mathbb{N}}$ denote a bounded sequence in $H^1_{\rho_*}$. Since $\mathcal{H}^{-1}$ is a bounded operator, also the sequence $\{\psi_n\}_{n\in\mathbb{N}}$ with $\mathcal{H}^{-1}\xi_n = \psi_n$ is bounded. Hence, because $H^1_{\rho_*}$ is a separable Hilbert space, there exist ξ and ψ in $H^1_{\rho_*}$ and subsequences (that we do not relabel) such that $\{\xi_n\}_{n\in\mathbb{N}}$ and $\{\psi_n\}_{n\in\mathbb{N}}$ converge weekly to ξ and ψ, respectively. By uniqueness of the weak limit, it is $\mathcal{H}^{-1}\xi = \psi$. Moreover, by Lemma 4 in the appendix, this convergence is strong in $L^2(B_1, \rho_*^{2-m}dx)$. Integrating by parts and using the definition of ψ_n, we have

$$\int \rho_*|\nabla\psi_n|^2\,dx \;=\; \frac{1}{m}\int \rho_*^{2-m}\psi_n\xi_n\,dx.$$

By the strong convergence in $L^2(B_1, \rho_*^{2-m}dx)$ we conclude that

$$\lim_{n\uparrow\infty}\frac{1}{m}\int \rho_*^{2-m}\psi_n\xi_n\,dx \;=\; \frac{1}{m}\int \rho_*^{2-m}\psi\xi\,dx \;=\; \int \rho_*|\nabla\psi|^2\,dx,$$

and the last identity follows from integrating by parts again and using $\mathcal{H}^{-1}\xi = \psi$. This shows that the $H^1_{\rho_*}$ norms of the sequence $\{\psi_n\}_{n\in\mathbb{N}}$ converge, and together with weak convergence, this yields convergence of $\{\psi_n\}_{n\in\mathbb{N}}$ (or equivalently of $\{\mathcal{H}^{-1}\xi_n\}_{n\in\mathbb{N}}$) in $H^1_{\rho_*}$. We deduce that $\mathcal{H}^{-1}$ is compact. □

2.3 Separation of variables in spherical coordinates

In this subsection, we most widely summarize Subsection 4.1 of [9]. As our argumentation for the porous medium equation follows closely the one for the fast-diffusion equation by Denzler & McCann, we entirely skip proofs and refer to [9] for details. An important step in the work of Denzler and McCann is the change of perspective in the spectral analysis of the displacement Hessian $\mathcal{H}$ which comes

along with the transformation of the operator into spherical coordinates. In fact, the change of variables into spherical coordinates is motivated by the crucial insight that the displacement Hessian $\mathcal{H}$ and the spherical Laplacian $\Delta_{\mathbb{S}^{N-1}}$ (i.e., the Laplace–Beltrami operator on the sphere $\mathbb{S}^{N-1}$) commute. This is basically a consequence of the fact that the evolution commutes with rotations. Consequently, both operators can be simultaneously diagonalized. In particular, as the spectrum of $\Delta_{\mathbb{S}^{N-1}}$ is well-known, our spectral analysis will reduce to the study of the radial part of $\mathcal{H}$, which amounts to a one-dimensional operator. We remark that this Ansatz is frequently used in the spectral analysis of Schrödinger operators, cf. [21].

We transform $x \in \mathbb{R}^N$ into spherical coordinates, i.e., $x = r\omega$ with $(r, \omega) \in [0, \infty) \times \mathbb{S}^{N-1}$, and recall that under this transformation, the Laplacian reads

$$\Delta_{\mathbb{R}^N} \;=\; \partial_r^2 + \frac{N-1}{r}\partial_r + \frac{1}{r^2}\Delta_{\mathbb{S}^{N-1}}.$$

Here $\Delta_{\mathbb{S}^{N-1}}$ denotes the spherical Laplacian. Since the Barenblatt profile ρ_* is a radially symmetric function, i.e., $\rho_* = \rho_*(r)$, one readily checks that $\mathcal{H}$ and $\Delta_{\mathbb{S}^{N-1}}$ commute, cf. (14).

The eigenvalues of the spherical Laplacian $\Delta_{\mathbb{S}^{N-1}}$ are $\mu_\ell = \ell(\ell + N - 2)$ with multiplicity $M_\ell = \frac{(N+\ell-3)!(N+2\ell-2)}{\ell!(N-2)!}$ where $M_0 = 1$. That is,

$$-\Delta_{\mathbb{S}^{N-1}} Y_{\ell m} \;=\; \mu_\ell Y_{\ell m} \quad \text{for } \ell \in \mathbb{N} \cup \{0\} \text{ and } m \in \{1, \dots, M_\ell\},$$

and the eigenfunctions $Y_{\ell m}$ are the spherical harmonics $Y_{\ell m} : \mathbb{S}^{N-1} \to \mathbb{R}$, which form a complete orthonormal basis for $L^2(\mathbb{S}^{N-1}, d\omega)$:

$$\int_{\mathbb{S}^{N-1}} Y_{\ell m}(\omega) Y_{\ell' m'}(\omega)\, d\omega \;=\; \delta_{\ell\ell'}\delta_{mm'}. \tag{19}$$

For more details about the spectrum and the eigenfunctions of the spherical Laplacian, see also [13, Ch. 3].

Exploiting the orthonormality of the spherical harmonics, we observe that for every radially symmetric function $f : (0, 1) \to \mathbb{R}$ the quantity $\|fY_{\ell,m}\|_{H^1_{\rho_*}}$ is independent of the particular choice of $m \in \{1, \dots, M_\ell\}$,

$$\|fY_{\ell m}\|^2_{H^1_{\rho_*}} \;=\; \int_0^1 \left((f'(r))^2 + \frac{\mu_\ell}{r^2}(f(r))^2 \right) \rho_*(r) r^{N-1}\, dr.$$

This leads us to the definition of the norm $\|f\|_{H^1_\ell} = \|fY_{\ell m}\|_{H^1_{\rho_*}}$ and the corresponding Sobolev space

$$H^1_\ell \;=\; \left\{ f \in H^1_{loc}(0, 1) :\; \|f\|_{H^1_\ell} < \infty \right\}.$$

In the case $\ell = 0$ (i.e., $\mu_\ell = 0$), we have to identify functions that only differ by a constant. In fact, H^1_ℓ is a Hilbert space and there is an isometric Hilbert space isomorphism

$$H^1_{\rho_*} = \bigoplus_{\ell=0}^{\infty} \bigoplus_{m=1}^{M_\ell} H^1_\ell$$

given by

$$\begin{aligned}\psi(r\omega) &= \sum_{\ell=0}^{\infty}\sum_{m=1}^{M_\ell} f_{\ell m}(r)Y_{\ell m}(\omega),\\ f_{\ell m}(r) &= \int_{\mathbb{S}^{N-1}} \psi(r\omega)Y_{\ell m}(\omega)\,d\omega,\end{aligned}$$

where the series are converging in $L^2(\mathbb{S}^{N-1}, d\omega)$, and the isometry reads

$$\|\psi\|^2_{H^1_{\rho_*}} = \sum_{l=0}^{\infty}\sum_{m=1}^{M_\ell} \|f_{\ell m}\|^2_{H^1_\ell}.$$

The orthogonal decomposition of the Hilbert space $H^1_{\rho_*}$ into eigenspaces generated by the spherical harmonics permits us to expand the displacement Hessian into a series of radially symmetric operators. More precisely, choosing $\psi \in \mathcal{D}(\mathcal{H})$ with $\psi(r\omega) = f(r)Y_{\ell m}(\omega)$, we have $\mathcal{H}\psi = (\mathcal{H}_\ell f)Y_{\ell m}$, where $\mathcal{H}_\ell$ is the orthogonal projection of $\mathcal{H}$ onto H^1_ℓ, namely

$$\left(\mathcal{H}_\ell f\right)(r) = -m\rho_*^{m-1}\left(f''(r) + \frac{N-1}{r}f'(r) - \frac{\mu_\ell}{r^2}f(r)\right) + rf'(r). \tag{20}$$

It is clear that this operator is again non-negative and self-adjoint with domain $\mathcal{D}(\mathcal{H}_\ell) = \{f \in H^1_\ell : fY_{\ell m} \in \mathcal{D}(\mathcal{H})\}$. For an explicit definition of $\mathcal{D}(\mathcal{H}_\ell)$, we have to investigate how the asymptotic boundary condition $\rho_*\nabla\psi\cdot\nu = 0$ behaves under the transformation into spherical coordinates. (This investigation is not necessary in the work of Denzler and McCann as they consider functions that are spread all over $\mathbb{R}^N$. In particular, in the fast-diffusion case the Barenblatt profile is positive everywhere.) The integrability condition on the free boundary ∂B_1 will become a selection criterion for identifying eigenfunctions of $\mathcal{H}_\ell$, and therefore, it has to be studied with care. For $\psi(r\omega) = f(r)Y_{\ell m}(\omega)$ and $\xi \in H^1_{\rho_*}$, we first compute that

$$-\int \nabla\cdot(\rho_*\nabla\psi)\,dx = -\int_0^1 \left(\rho_* f' r^{N-1}\right)' g_{\ell m}\,dr + \mu_\ell\int_0^1 \rho_* f g_{\ell m} r^{N-3}\,dr,$$

and

$$\int \rho_*\nabla\psi\cdot\nabla\xi\,dx = \int_0^1 \rho_*\left(f'g'_{\ell m} + \frac{\mu_\ell}{r^2}fg_{\ell m}\right)r^{N-1}\,dr,$$

where $g_{\ell m}(r) = \int_{\mathbb{S}^{N-1}}\xi(r\omega)Y_{\ell m}(\omega)\,d\omega$. Consequently, (17) becomes

$$\int_0^1 \rho_* f'g' r^{N-1}\,dr = -\int_0^1 (\rho_* f' r^{N-1})' g\,dr \quad \text{for all } g \in H^1_\ell. \tag{21}$$

As the boundary term at $r = 0$ automatically vanishes for every $f, g \in H^1_\ell$, we simply write $\rho_* f'|_{r=1} = 0$ to indicate that (21) holds. Now the domain of self-adjointness of $\mathcal{H}_\ell$ can be written as

$$\mathcal{D}(\mathcal{H}_\ell) = \left\{f \in H^3_{loc}(0,1)\cap H^1_\ell \;\middle|\; \mathcal{H}_\ell f \in H^1_\ell,\ \rho_* f'|_{r=1} = 0\right\}.$$

Remark 2. *In analogy to the statement* (18) *in Remark 1, we like to point out that the asymptotic boundary condition in the radially symmetric variables reduces to*

$$\lim_{r\uparrow 1} \rho_*(r) f'(r) \;=\; 0, \tag{22}$$

for every function $f \in C^\infty[0,1]$.

2.4 The eigenvalue problem for $\mathcal{H}_\ell$

Under the assumption that $\mathcal{H}$ has countably many eigenvalues accumulating at infinity, we have shown in Proposition 2 above that the spectrum of $\mathcal{H}$ is purely discrete and that the eigenfunctions form an orthonormal basis of $H^1_{\rho_*}$. According to the isomorphy established in the previous subsection, it is enough to remove this assumption for the radial components of $\mathcal{H}$. In the following, we solve the eigenvalue problem for the self-adjoint operator $\mathcal{H}_\ell$.

Proposition 3 (Radial eigenvalue problem)**.** *For every* $\ell \in \mathbb{N}$*, the eigenvalue problem*

$$\mathcal{H}_\ell f \;=\; \lambda f$$

in $\mathcal{D}(\mathcal{H}_\ell)$ *has exactly the eigenvalues*

$$\lambda_{\ell,k} \;=\; \ell + 2k + 2k(k+\ell+\frac{N}{2}-1)(m-1),$$

where $k \in \mathbb{N}_0$ *such that* $(\ell,k) \neq (0,0)$*. The corresponding eigenfunctions are the polynomials*

$$f_{\ell,k}(r) = r^\ell F(-k, \frac{1}{m-1} + \ell + \frac{N}{2} - 1 + k; \ell + \frac{N}{2}; r^2),$$

where $F(-k,b;c;z) = 1 + \sum_{n=1}^k \frac{(-k)_n (b)_n}{(c)_n n!} z^n$ *with* $(s)_n = s(s+1)\dots(s+n-1)$.

From the study of the operator $\mathcal{H}$ in Proposition 1 we already know that any eigenvalue of $\mathcal{H}_\ell$ must be real and positive, i.e., $\lambda > 0$. Notice that in view of the explicit formulas (20) & (4) for $\mathcal{H}_\ell$ and ρ_*, the equation $\mathcal{H}_\ell f = \lambda f$ reads

$$f'' + \left(\frac{N-1}{r} - \frac{2}{m-1}\frac{r}{1-r^2}\right) f' + \left(\frac{2\lambda}{m-1}\frac{1}{1-r^2} - \frac{\mu_\ell}{r^2}\right) f \;=\; 0 \quad \text{in } (0,1). \tag{23}$$

This is a linear differential equation of second order with singularities at the endpoints of the interval $(0,1)$. It is of Fuchsian type, that means, all singular points, here 0, 1, and ∞, are regular. For every given λ, this differential equation has two linearly independent families of solutions. It is well known that the study of second-order Fuchsian ODEs with three regular singular points is intertwined with the study of hypergeometric functions $F(a,b;c;z)$ defined by

$$F(a,b;c;z) \;=\; \sum_{n=0}^\infty \frac{(a)_n (b)_n}{(c)_n n!} z^n, \tag{24}$$

where $a, b, c, z \in \mathbb{R}$ and c is not a non-positive integer. Here, the notation involves the Pochhammer symbols (or extended factorials)

$$(s)_n = s(s+1)\dots(s+n-1), \quad \text{for } n \geq 1, \quad \text{and} \quad (s)_0 = 1.$$

It is easily verified that the series converges if $|z| < 1$. In the case that moreover $c > b > 0$, the hypergeometric function $F(a, b; c; z)$ has the integral representation

$$F(a,b;c;z) \;=\; \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^\infty t^{b-1}(1+t)^{a-c}(1+t-zt)^{-a}\, dt,$$

were $\Gamma(s)$ denotes Euler's gamma function. We finally quote the fact that is most relevant for our purposes: For every choice of $a, b, c, z \in \mathbb{R}$, c not a non-positive integer, the function $F(a, b; c; z)$ satisfies the *Hypergeometric differential equation*

$$F'' + \left(\frac{c}{z} - \frac{a+b-c+1}{1-z}\right) F' - \frac{ab}{z(1-z)} F \;=\; 0 \quad \text{in } (0,1), \tag{25}$$

where the primes indicate derivatives with respect to z, i.e, $F' = \partial_z F$ and $F'' = \partial_z^2 F$. For detailed discussions of hypergeometric functions, we refer to [25, 26, 4]. A compact catalogue of the most important facts can be found in [1].

Now, we come back to the differential equation (23) and its relation to the hypergeometric equation (25). Our goal is to transform (23) into the hypergeometric differential equation (25), for which all solutions and their asymptotics at the singular points 0, 1, and ∞ are well-known. Making the Ansatz $f(r) = r^\gamma F(a, b; c; r^2)$ and supposing that f is a solution, (23) transforms into

$$\begin{aligned} F'' &+ \left(\frac{2\gamma+N}{2}\frac{1}{z} - \frac{1}{m-1}\frac{1}{1-z}\right) F' \\ &+ \left(\frac{\gamma(\gamma+N-2)-\mu_\ell}{4}\frac{1}{z^2} - \frac{\gamma-\lambda}{2(m-1)}\frac{1}{z(1-z)}\right) F \;=\; 0, \end{aligned} \tag{26}$$

with $z = r^2$. Comparing this differential equation with (25) and identifying a, b, c, and d yields a first solution to (23). A second linearly independent solution can be deduced from the well-known (but complex) theory of hypergeometric functions. We have the following

Lemma 1. *Let*

$$a \;=\; \frac{1}{2}\left(\frac{1}{m-1} + \ell + \frac{N}{2} - 1\right) - \frac{1}{2}\sqrt{q}, \tag{27}$$

$$b \;=\; \frac{1}{2}\left(\frac{1}{m-1} + \ell + \frac{N}{2} - 1\right) + \frac{1}{2}\sqrt{q}, \tag{28}$$

$$q \;=\; \frac{2\lambda}{m-1} + \left(\ell + \frac{N}{2} - 1\right)^2 + \frac{N-2}{m-1} + \frac{1}{(m-1)^2}, \tag{29}$$

$$c \;=\; \ell + \frac{N}{2}. \tag{30}$$

Then the differential equation (23) *has two linearly independent solutions* $f_1(r)$ *and* $f_2(r)$*. A first solution is of the form*

$$f_1(r) \;=\; r^\ell F(a,b;c;r^2),$$

where F *denotes the hypergeometric function defined in* (24)*. In the case where* c *is not a positive integer, a second solution* f_2 *is given by*

$$f_2(r) \;=\; r^{2-\ell-N} F(a+1-c, b+1-c; 2-c; r^2).$$

If c *is a positive integer,* $c = k \in \mathbb{N}$*, then a second solution has the asymptotics*

$$f_2(r) \;\sim\; \left\{ \begin{array}{ll} \ln r & \text{for } k=1 \\ r^{\ell+2-2k} & \text{for } k \geq 2 \end{array} \right\} \quad \text{as } 0 < r \ll 1,$$

and

$$f_2'(r) \;\sim\; r^{\ell+1-2k} \quad \text{as } 0 < r \ll 1.$$

As the second solution f_2 will be discarded as a potential eigenfunction of the eigenvalue problem (23) in the proof of Proposition 3 below, it suffices at this point to quote only its asymptotic behavior at zero for positive integers c.

Proof of Lemma 1. A first solution to (23) can be obtained by identifying a set of values γ, a, b, and c for which (26) turns into the form (25), and setting $f_1(r) = r^\gamma F(a,b;c;r^2)$. The form of a second linearly independent solution depends on the particular values of a, b, and c. In the simplest case, the second solution is obtained by an appropriate change of the dependent and independent variables. In some cases, however, the construction of the second solution relies deeply on the theory of hypergeometric functions, and we have to refer to the relevant literature.

We start identifying values for γ, a, b, and c such that the hypergeometric function $F(a,b;c;z)$ solves (26). Comparing (26) and (25), we immediately see that $\gamma(\gamma + N - 2) = \mu_\ell$. In view of the definition $\mu_\ell = \ell(\ell+N-2)$ this enforces either $\gamma = \ell$ or $\gamma = 2 - N - \ell$. We choose $\gamma = \ell$. Moreover, the constant c is determined by $\gamma(=\ell)$ and N only, namely by $c = \ell + \frac{N}{2}$, i.e., (30). Finally, the values for a and b can be derived from the identities $a + b - c + 1 = \frac{1}{m-1}$ and $ab = \frac{\ell-\lambda}{2(m-1)}$. A short computation yields the two solutions $a_\pm = b_\mp = \frac{1}{2}\left(\frac{1}{m-1} + \ell + \frac{N}{2} - 1\right) \pm \frac{1}{2}\sqrt{q}$, where $q = \frac{2\lambda}{m-1} + (\ell + \frac{N}{2} - 1)^2 + \frac{N-2}{m-1} + \frac{1}{(m-1)^2}$, and this quantity is positive since λ is positive. We choose $a = a_- = b_+$ and $b = a_+ = b_-$, i.e., (27)–(29). Observe that c is not a non-positive integer since $N \geq 2$ and $\ell \geq 0$, so that $F(a,b;c;z)$ is well defined for a, b, and c as above.

Depending on the particular values of a, b and c, we will in the following compute a second linearly independent solution of (23). This solution will again be of the form $f_2(r) = r^\ell F(\tilde{a},\tilde{b};\tilde{c};r^2)$, where $\tilde{F} := F(\tilde{a},\tilde{b};\tilde{c};r^2)$ is a second linearly independent

solution of the hypergeometric equation (25). Indeed, computing the Wronskian $W(f_1, f_2)$ yields

$$\begin{aligned} W(f_1, f_2) &= f_1' f_2 - f_1 f_2' \\ &= 2r^{2\ell+1}(F'\tilde{F} - F\tilde{F}') \\ &= 2r^{2\ell+1} W(F, \tilde{F}) \neq 0 \end{aligned}$$

since F and $\tilde{F}$ are linearly independent.

In our derivation of $\tilde{F}$, we follow [4, Ch. 8]. If c is not a positive integer, then a second linearly independent solution to the hypergeometric differential equation (25) is given by $z^{1-c}F(a+1-c, b+1-c; 2-c; z)$ (cf. [4, eq. (8.2.6)]) and thus $f_2(r) = r^{2-\ell-N}F(a+1-c, b+1-c; 2-c; r^2)$ is a second linearly independent solution to (23). Notice that f_1 coincides with f_2 in the case $c = 1$ and the latter function is not even defined for larger integer values of c. For integers $c = k \geq 1$, a second solution to the hypergeometric differential equation is given by

$$\begin{aligned} G(a, b; k; z) &= \frac{(\log z) F(a, b; k; z)}{\Gamma(a+1-k)\Gamma(b+1-k)(k-1)!} \\ &+ \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(k)_n n!} \frac{\theta(a+n) + \theta(b+n) - \theta(n+1) - \theta(k+n)}{\Gamma(a+1-k)\Gamma(b+1-k)(k-1)!} z^n \\ &+ \frac{(-1)^k (k-2)!}{\Gamma(a)\Gamma(b)} \sum_{n=0}^{k-2} \frac{(a+1-k)_n (b+1-k)_n}{(2-k)_n n!} z^{n+1-k}, \end{aligned}$$

where $\theta(s) = \frac{\Gamma'(s)}{\Gamma(s)}$ and the convention that the last term is zero if $k = 1$, cf. [4, eq. (8.4.4)]. We remark that $F(a, b; k; 0) = 1$. Inspecting the asymptotic behavior at the origin and eventually redefining $G(a, b; k; z)$ by multiplying by $\Gamma(a)$ and/or $\Gamma(b)$ if a and/or b is a nonpositive integer, we see that

$$G(a, b; k; z) \sim \left\{ \begin{array}{ll} \ln z & \text{for } k = 1 \\ z^{1-k} & \text{for } k \geq 2 \end{array} \right\} \quad \text{as } 0 < z \ll 1.$$

and

$$\partial_z G(a, b; k; z) \sim z^{-k} \quad \text{as } 0 < z \ll 1.$$

(See also the discussion on page 275 in [4].) It remains to set $f_2(r) = r^{\ell} G(a, b; k; r^2)$ and notice that $k = 1$ precisely if $\ell = 0$ and $N = 2$. This proves Lemma 1. □

We are now in the position to present the

Proof of Proposition 3. We consider the eigenvalue problem $\mathcal{H}_\ell f = \lambda f$ for $f \in \mathcal{D}(\mathcal{H}_\ell)$. By the discussion in the previous section and Proposition 1, we already know that every eigenvalue λ must be positive. It is readily checked that the eigenvalue equation for $\mathcal{H}_\ell$ is equivalent to the differential equation (23). This equation is solvable for every $\lambda > 0$ by Lemma 1, and a solution f to (23) is an eigenfunction

of $\mathcal{H}_\ell$ if and only if $f \in \mathcal{D}(\mathcal{H}_\ell)$. Moreover, f can be written as a linear combination of the linearly independent solutions f_1 and f_2. As solutions of (23) are automatically smooth in $(0,1)$ by interior regularity results, we have to study the asymptotic behavior of f_1 and f_2 at the singular points 0 and 1 in order to decide whether $f_1, f_2 \in \mathcal{D}(\mathcal{H}_\ell)$.

For further references we quote that

$$\partial_z F(a,b;c;z) \;=\; \frac{ab}{c} F(a+1,b+1;c+1;z),$$

cf. [4, eq. (8.2.3)], and therefore

$$f_1'(r) \;=\; \ell r^{\ell-1} F(a,b;c;r^2) + 2\frac{ab}{c} r^{\ell+1} F(a+1,b+1;c+1;r^2), \tag{31}$$

and, provided c is not a positive integer,

$$\begin{aligned} f_2'(r) \;&=\; (2-\ell-N) r^{1-\ell-N} F(a-c+1,b-c+1,2-c;r^2) \\ &\quad + 2\frac{(a-c+1)(b-c+1)}{2-c} r^{3-\ell-N} F(a-c+2,b-c+2;3-c;r^2). \end{aligned} \tag{32}$$

Moreover, one easily computes that

$$F(a,b;c;0) \;=\; 1 \tag{33}$$

if c is not a non-positive integer.

Step 1. It holds $f_2 \notin H^1_\ell$, and thus f_2 is not an eigenfunction of $\mathcal{H}_\ell$.
We first consider the case where $c = k$ is an integer. Then, by Lemma 1 it is $f_2'(r) \sim r^{\ell+1-2k}$ asymptotically at the origin, and thus, using $2k = 2\ell + N$ by (30)

$$\rho_*(r)(f_2'(r))^2 r^{N-1} \;\sim\; r^{2\ell+1-4k+N} \;=\; r^{1-2k} \quad \text{as } 0 < r \ll 1.$$

The term on the right is not integrable for any $k \geq 1$, and thus $\|f_2\|_{H^1_\ell} = \infty$. This shows that $f_2 \notin H^1_\ell$ in the case where $c = k$ is an integer. If c is not an integer, then our argument is based on (32) and (33). In fact, denoting by $C_{a,b,c}$ a constant that depends only on a, b, and c, we have $f_2'(r) \approx (2-\ell-N) r^{1-\ell-N} + C_{a,b,c} r^{3-\ell-N} \sim r^{1-\ell-N}$ asymptotically at the origin. Here we have used that $\ell + N \neq 2$ because $c \neq 1$. Therefore

$$\rho_*(r)(f_2'(r))^2 r^{N-1} \;\sim\; r^{1-2\ell-N} \;=\; r^{1-2c} \quad \text{as } 0 < r \ll 1.$$

Again, the right hand side is not integrable since $c > 1$ and thus $f_2 \notin H^1_\ell$ in the case where c is not an integer.

Step 2. For $-a \notin \mathbb{N} \cup \{0\}$, it holds $f_1 \notin \mathcal{D}(\mathcal{H}_\ell)$, and thus f_1 is not an eigenfunction of $\mathcal{H}_\ell$.
To rule out solutions f_1 as eigenfunctions of $\mathcal{H}_\ell$ in the case $-a \notin \mathbb{N} \cup \{0\}$, we investigate the asymptotics at the singularity $r = 1$. Since solutions f_1 are smooth in $(0,1)$, the asymptotic boundary conditions in the definition of $\mathcal{D}(\mathcal{H}_\ell)$ can be explicitly

checked via (22). In fact, eigenfunctions must necessarily satisfy $\lim_{r\uparrow 1} \rho_*(r) f'(r) = 0$.

In the following, we show that this condition fails if $-a \notin \mathbb{N} \cup \{0\}$. In the case that $a+b-c \notin \mathbb{Z}$, we have the linear transformation formula

$$\begin{aligned} F(a,b;c;z) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F(a,b;a+b-c+1;1-z) \\ &\quad + (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} F(c-a,c-b;c-a-b+1;1-z), \end{aligned}$$

cf. [1, 15.3.6]. In view of (33) and $a+b-c \notin \mathbb{Z}$, it is

$$F(a,b;c;z) \approx \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} + (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} \quad \text{as } 0 < 1-z \ll 1.$$

Invoking (31), $b > 0$, and $a+b-c+1 = \frac{1}{m-1}$, we deduce that

$$\rho_*(r) f_1'(r) \sim 2 \frac{\Gamma(c)\Gamma(a+b-c+1)}{\Gamma(a)\Gamma(b)} (1-r^2)^{\frac{1}{m-1}+c-a-b-1} \sim 1 \quad \text{as } 0 < 1-r \ll 1,$$

and thus (22) is violated. Consequently, f_1 is not in $\mathcal{D}(\mathcal{H}_\ell)$, and thus not an eigenfunction of $\mathcal{H}_\ell$.

As the above linear transformation has a pole when $a+b-c \in \mathbb{Z}$, we have to use substitute formulas to investigate the limiting behavior at $z=1$. Such transformation formulas can be found in [1, 15.3.10, 15.3.12]. Instead of displaying the explicit expressions in the sequel, we will just quote the limiting behavior of $F(a,b;c;z)$ as z converges to 1. We note that $c < a+b+1$ since $a+b-c+1 = \frac{1}{m-1} > 0$. We have

$$F(a,b;a+b;z) \approx -\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \ln(1-z) \quad \text{as } 0 < 1-z \ll 1,$$

and for $k \in \mathbb{N}$,

$$F(a,b;a+b-k;z) \approx \frac{\Gamma(k)\Gamma(a+b-k)}{\Gamma(a)\Gamma(b)} (1-z)^{-k} \quad \text{as } 0 < 1-z \ll 1.$$

For every $k \in \mathbb{N} \cup \{0\}$, the case $a+b-c=k$ is equivalent to $\frac{1}{m-1} = k+1$, and thus, a short computation using (31) yields

$$\rho_*(r) f_1'(r) \sim (1-r^2)^{\frac{1}{m-1}-(k+1)} = 1 \quad \text{as } 0 < 1-r \ll 1.$$

This violates (22) and thus f_1 is not an eigenfunction of $\mathcal{H}_\ell$.

Step 3. For $-a \in \mathbb{N} \cup \{0\}$, it holds $f_1 \in \mathcal{D}(\mathcal{H}_\ell)$, and thus f_1 is an eigenfunction of $\mathcal{H}_\ell$. The corresponding eigenvalue is $\lambda = \ell + 2k + 2k\left(k+\ell+\frac{N}{2}-1\right)(m-1)$.

We set $a = -k$ for some $k \in \mathbb{N} \cup \{0\}$ and observe that the hypergeometric series in (24) terminates, namely

$$F(-k, b; c; z) = \sum_{n=0}^{k} \frac{(a)_n (b)_n}{(c)_n n!} z^n.$$

It follows that f_1 is a polynomial of degree $2k + \ell$ and thus a continuous function on $[0, 1]$. In particular, the asymptotic boundary condition (22) is trivially satisfied. Regarding the integrability of f_1, we only have study the singularity 0 if $\ell \geq 1$. In this case, we have that $(f_1(r))^2 r^{N-3} \approx r^{2\ell+N-3}$, which is integrable, and thus $f_1 \in H^1_\ell$. Moreover, since $\mathcal{H}_\ell f_1 = \lambda f_1$ we also know that $\mathcal{H}_\ell f_1 \in H^1_\ell$, and thus $f_1 \in \mathcal{D}(\mathcal{H}_\ell)$. We conclude that f_1 is an eigenfunction of $\mathcal{H}_\ell$ if a is a nonpositive integer.

It remains to express the eigenvalue λ in terms the fixed variables. A straight forward computation based on the expression (27) yields that

$$\lambda = \ell + 2k + 2k \left(k + \ell + \frac{N}{2} - 1 \right) (m - 1).$$

This concludes the proof of Proposition 3. □

Appendix: Elliptic theory in $H^1_{\rho_*}$

This appendix provides some results related to elliptic problems in $H^1_{\rho_*}$ that apply in the discussion of the displacement Hessian $\mathcal{H}$ in Section 2.2. The statements in the Lemmas 2–5 below are the analogs of very classical facts in standard Sobolev theory: density of smooth functions, a Hardy–Poincaré inequality, a Rellich lemma, the Poisson problem. Our Hilbert space $H^1_{\rho_*}$ differs from the well-studied Sobolev space $H^1(B_1)$ only by the weight ρ_*, which is, by the regularity of the Barenblatt profile in the ball B_1, a very mild variation: ρ_* is finite and degenerates only on the boundary ∂B_1. In this regard, it is not surprising that the results in $H^1_{\rho_*}$ can be proved analogously to or based on the known theory.

We first give an overview of the main facts. We start with a classical density statement.

Lemma 2 (Density of smooth functions)**.** *For every $\psi \in H^1_{\rho_*}$, there exists a sequence $\{\psi_\nu\}_{\nu \in \mathbb{N}}$ in $C^\infty(\bar{B}_1)$ such that*

$$\lim_{\nu \uparrow \infty} \int \rho_* |\nabla(\psi - \psi_\nu)|^2 \, dx \;=\; 0.$$

The next result is an embedding theorem between weighted Lebesgue and Sobolev spaces, cf. [9, Corollary 13].

Lemma 3 (Hardy–Poincaré inequality)**.** *Let $p > 1$ such that $p + m \geq 3$. Then*

$$\inf_{c\in\mathbb{R}} \int \rho_*^{p-m}(\psi - c)^2\,dx \;\lesssim\; \int \rho_*|\nabla\psi|^2\,dx \tag{34}$$

for any $\psi \in H^1_{\rho_*}$.

Notice that the infimum in the statement of the Hardy–Poincaré inequality is attained by $c = \int \rho_*^{p-m}\psi\,dx / \int \rho_*^{p-m}\,dx$. The inequality is sharp if $p+m = 3$. In most cases, we will apply (34) with $p = 2$. The general result is used in the proof of the following compactness result.

Lemma 4 (Rellich lemma)**.** *The embedding of $H^1_{\rho_*}$ in $L^2(B_1, \rho_*^{2-m}dx)$ is compact.*

We finally study the Poisson problem in $H^1_{\rho_*}$.

Lemma 5 (Poisson problem)**.** *For every $u \in L^2(B_1, \rho_*^{m-2}dx)$ with $\int u\,dx = 0$ there exists a unique (up to an additive constant) $\psi \in H^2_{\mathrm{loc}}(B_1) \cap H^1_{\rho_*}$ such that*

$$\begin{aligned} -\nabla\cdot(\rho_*\nabla\psi) &= u \quad \text{in } B_1 \\ \rho_*\nabla\psi\cdot\nu &= 0 \quad \text{on } \partial B_1. \end{aligned}$$

As before, we understand the boundary conditions in the above lemma in the sense of (17).

We first address the proof of Lemma 2, which uses classical approximation techniques, see e.g., [11, Sec. 5.3].

Proof of Lemma 2. We fix $\psi \in H^1_{\rho_*}$ and notice that that $\psi \in H^1(K)$ for every compact set $K \subset B_1$ since ρ_* is positive and finite away from ∂B_1. In a first step, we show that ψ can be approximated by $C^\infty(B_1)$ functions. For this purpose, we consider $A_0 = B_{\frac{2}{3}}$ and annuli $A_\nu = B_{\frac{\nu+2}{\nu+3}} \setminus B_{\frac{\nu}{\nu+1}}$ for every $\nu \in \mathbb{N}$, where B_r denotes the ball of radius r around the origin, to the effect of $B_1 = \bigcup_{\nu=0}^\infty A_\nu$. Let $\{\eta_\nu\}_{\nu\in\mathbb{N}_0}$ be a partition of unity subordinate the covering $\{A_\nu\}_{\nu\in\mathbb{N}_0}$ and let $\{\zeta_\mu\}_{\mu\in\mathbb{N}_0}$ be a sequence of standard mollifiers. We fix $\varepsilon > 0$ arbitrarily. Since $\eta_\nu\psi \in H^1(A_\nu)$, for any $\nu \in \mathbb{N}_0$ there exists a $\mu = \mu(\varepsilon,\nu) \in \mathbb{N}_0$ such that

$$\left(\int \rho_*|\nabla\left(\eta_\nu\psi - (\eta_\nu\psi)*\zeta_\mu\right)|^2\,dx\right)^{1/2} \;\leq\; \frac{\varepsilon}{2^{\nu+1}}$$

and

$$\operatorname{supp}\left((\eta_\nu\psi)*\zeta_\mu\right) \subset B_{\frac{\nu+3}{\nu+4}} \setminus B_{\frac{\nu-1}{\nu}}.$$

We define $\tilde\psi = \sum_{\nu=0}^\infty \psi_\nu$ where $\psi_\nu = (\eta_\nu\psi)*\zeta_\mu$. We obviously have $\tilde\psi \in C^\infty(B_1)$. Moreover, since $\psi = \sum_{\nu=0}^\infty \eta_\nu\psi$, we have for every compact set $K \subset B_1$:

$$\begin{aligned} \left(\int_K \rho_*|\nabla(\psi-\tilde\psi)|^2\,dx\right)^{1/2} &\leq \sum_{\nu=0}^\infty \left(\int_K \rho_*|\nabla(\eta_\nu\psi - \psi_\nu)|^2\,dx\right)^{1/2} \\ &\leq \sum_{\nu=0}^\infty \frac{\varepsilon}{2^{\nu+1}} \;=\; \varepsilon. \end{aligned}$$

Taking the supremum over all K, this shows that $C^\infty(B_1)$ is dense in $H^1_{\rho_*}$.

In order to prove density of functions that are smooth up to the boundary, we may, due to the above argumentation, suppose that $\psi \in C^\infty(B_1)$. For $x \in B_1$ and $\nu \in \mathbb{N}$, we define $\psi_\nu(x) = \psi((1-\frac{1}{\nu})x)$. It follows that $\psi_\nu \in C^\infty(\bar{B}_1)$ and

$$\int \rho_* |\nabla(\psi - \psi_\nu)|^2 \, dx \ \overset{\nu\uparrow\infty}{\longrightarrow}\ 0$$

by the dominated convergence theorem. □

We now turn to the proof of Lemma 3, which relies on the following one-dimensional inequality:

Lemma 6 (Double-weight Hardy inequality)**.** *Let $p > 1$ such that $p + m \geq 3$. Then*

$$\inf_{c\in\mathbb{R}} \int_0^1 (1-r)^{\frac{p-m}{m-1}} (\psi - c)^2 r^{N-1} \, dr \ \lesssim\ \int_0^1 (1-r)^{\frac{1}{m-1}} (\psi')^2 r^{N-1} \, dr \tag{35}$$

for any $\psi \in C^1[0,1]$.

Notice that the above inequality is optimal for $p + m = 3$ in the case $m \neq 2$. For $p = 1$ and $m = 2$, however, this estimate fails (logarithmically) as the underlying Hardy inequality is critical.

Proof of Lemma 3. Since $C^\infty(\bar{B}_1)$ is dense in $H^1_{\rho_*}$ by Lemma (1), it is enough to consider the case where ψ is smooth up to the boundary. We prove a slightly stronger statement by choosing $c = \int \rho_*^{p-1} |x|^{-1} \psi \, dx / \int \rho_*^{p-1} |x|^{-1} \, dx$, or equivalently by assuming that

$$\int \rho_*^{p-1} |x|^{-1} \psi \, dx \ =\ 0. \tag{36}$$

Notice that the integral is well-defined for every $\psi \in C^\infty(\bar{B}_1)$ by the assumption on p and since $N \geq 2$. We rewrite the statement in spherical coordinates: Let $x = r\omega$ and $\hat{\psi}(r,\omega) = \psi(x)$ where $r = |x|$ and $\omega = x/|x|$. Then $|\nabla\psi|^2 = (\partial_r \hat{\psi})^2 + \frac{1}{r^2} |\nabla_{\mathbb{S}^{N-1}} \hat{\psi}|^2$, where $\nabla_{\mathbb{S}^{N-1}}$ denotes the tangential gradient and thus, by the definition of ρ_*, (34) follows from

$$\begin{aligned} &\int_0^1 \int_{\mathbb{S}^{N-1}} (1-r)^{\frac{p-m}{m-1}} \hat{\psi}^2 r^{N-1} \, d\omega dr \\ &\quad\lesssim\ \int_0^1 \int_{\mathbb{S}^{N-1}} (1-r)^{\frac{1}{m-1}} \left((\partial_r \hat{\psi})^2 + \frac{1}{r^2} |\nabla_{\mathbb{S}^{N-1}} \hat{\psi}|^2 \right) r^{N-1} \, d\omega dr. \end{aligned} \tag{37}$$

For any fixed ω, we consider $\psi_\omega(r) = \hat{\psi}(r,\omega)$. Since $\psi_\omega \in C^\infty[0,1]$, we can apply Lemma 6 componentwise for any $\omega \in \mathbb{S}^{N-1}$. Observe that the infimum in (35) is attained by a constant $c \sim \int_0^1 (1-r)^{\frac{p-m}{m-1}} \psi_\omega r^{N-1} \, dr$, so that (35) and integration over $\mathbb{S}^{N-1}$ yield

$$\begin{aligned} &\int_{\mathbb{S}^{N-1}} \int_0^1 (1-r)^{\frac{p-m}{m-1}} \hat{\psi}^2 r^{N-1} \, dr d\omega \\ &\quad\lesssim\ \int_{\mathbb{S}^{N-1}} \left(\int_0^1 (1-r)^{\frac{p-m}{m-1}} \hat{\psi} r^{N-1} \, dr \right)^2 d\omega + \int_{\mathbb{S}^{N-1}} \int_0^1 (1-r)^{\frac{1}{m-1}} (\partial_r \hat{\psi})^2 r^{N-1} \, dr d\omega. \end{aligned}$$

To derive (37), we need to control the first term on the right. Integrating by parts and applying the Cauchy–Schwarz inequality, we compute

$$\begin{aligned}
&\left|\int_0^1 (1-r)^{\frac{p-m}{m-1}}\hat\psi r^{N-1}\,dr\right| \\
&\quad= \left|\frac{m-1}{p-1}(N-1)\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr + \frac{m-1}{p-1}\int_0^1 (1-r)^{\frac{p-1}{m-1}}\partial_r\hat\psi r^{N-1}\,dr\right| \\
&\quad\lesssim \left|\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr\right| \\
&\qquad + \left(\int_0^1 (1-r)^{\frac{2p-3}{m-1}} r^{N-1}\,dr\right)^{1/2}\left(\int_0^1 (1-r)^{\frac{1}{m-1}}(\partial_r\hat\psi)^2 r^{N-1}\,dr\right)^{1/2}.
\end{aligned}$$

Now that the above estimate becomes

$$\begin{aligned}
&\int_{\mathbb{S}^{N-1}}\int_0^1 (1-r)^{\frac{p-m}{m-1}}(\hat\psi)^2 r^{N-1}\,dr d\omega \\
&\quad\lesssim \int_{\mathbb{S}^{N-1}}\left(\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr\right)^2 d\omega + \int_{\mathbb{S}^{N-1}}\int_0^1 (1-r)^{\frac{1}{m-1}}(\partial_r\hat\psi)^2 r^{N-1}\,dr d\omega,
\end{aligned}$$

and the statement in (37) follows from the estimate

$$\int_{\mathbb{S}^{N-1}}\left(\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr\right)^2 d\omega \;\lesssim\; \int_{\mathbb{S}^{N-1}}\int_0^1 (1-r)^{\frac{1}{m-1}}|\nabla_{\mathbb{S}^{N-1}}\hat\psi|^2 r^{N-3}\,dr d\omega. \tag{38}$$

To prove (38), we start by recalling the Poincaré inequality on a sphere (cf. [14, Theorem 2.10])

$$\int_{\mathbb{S}^{N-1}}\xi^2\,d\omega \;\lesssim \left(\int_{\mathbb{S}^{N-1}}\xi\,d\omega\right)^2 + \int_{\mathbb{S}^{N-1}}|\nabla_{\mathbb{S}^{N-1}}\xi|^2\,d\omega,$$

which we apply to $\xi = \int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr$ to the effect of

$$\begin{aligned}
&\int_{\mathbb{S}^{N-1}}\left(\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr\right)^2 d\omega \\
&\quad\lesssim \left(\int_{\mathbb{S}^{N-1}}\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr d\omega\right)^2 \\
&\qquad + \int_{\mathbb{S}^{N-1}}\left(\int_0^1 (1-r)^{\frac{p-1}{m-1}}|\nabla_{\mathbb{S}^{N-1}}\hat\psi| r^{N-2}\,dr\right)^2 d\omega.
\end{aligned}$$

The first term on the right vanishes since

$$\int_{\mathbb{S}^{N-1}}\int_0^1 (1-r)^{\frac{p-1}{m-1}}\hat\psi r^{N-2}\,dr d\omega \;\sim\; \int \rho_*^{p-1}|x|^{-1}\psi\,dx \overset{(36)}{=} 0.$$

The second term can be estimated using the Cauchy–Schwarz inequality in the inner integral:

$$\begin{aligned}&\int_{\mathbb{S}^{N-1}}\left(\int_0^1(1-r)^{\frac{p-1}{m-1}}|\nabla_{\mathbb{S}^{N-1}}\hat{\psi}|r^{N-2}\,dr\right)^2 d\omega\\ &\quad\le\quad\int_0^1(1-r)^{\frac{2p-3}{m-1}}r^{N-1}\,dr\int_{\mathbb{S}^{N-1}}\int_0^1(1-r)^{\frac{1}{m-1}}|\nabla_{\mathbb{S}^{N-1}}\hat{\psi}|^2r^{N-3}\,dr d\omega.\end{aligned}$$

Since the prefactor is bounded by the assumption on p, this proves (38). □

Proof of Lemma 6. The statement (35) follows as a combination of the two Hardy inequalities

$$\int_0^{1/2}(\psi-\psi(1/2))^2r^{N-1}\,dr\;\lesssim\;\int_0^{1/2}(\psi')^2r^{N+1}\,dr \tag{39}$$

and

$$\int_{1/2}^1(1-r)^{\frac{p-m}{m-1}}(\psi-\psi(1/2))^2\,dr\;\lesssim\;\int_{1/2}^1(1-r)^{\frac{1}{m-1}}(\psi')^2\,dr. \tag{40}$$

Indeed,

$$\begin{aligned}&\int_0^1(1-r)^{\frac{p-m}{m-1}}(\psi-\psi(1/2))^2r^{N-1}\,dr\\ &\quad\lesssim\quad\int_0^{1/2}(\psi-\psi(1/2))^2r^{N-1}\,dr+\int_{1/2}^1(1-r)^{\frac{p-m}{m-1}}(\psi-\psi(1/2))^2\,dr\\ &\quad\overset{(39)\&(40)}{\lesssim}\quad\int_0^{1/2}(\psi')^2r^{N+1}\,dr+\int_{1/2}^1(1-r)^{\frac{1}{m-1}}(\psi')^2\,dr\\ &\quad\lesssim\quad\int_0^1(1-r)^{\frac{1}{m-1}}(\psi')^2r^{N-1}\,dr,\end{aligned}$$

and the term on the left can be bounded from below by taking the infimum over all $c=\psi(1/2)\in\mathbb{R}$.

For proving (39) and (40), it is enough to consider the case where $\psi(1/2)=0$. The proof of the Hardy inequality (39) is standard:

$$\begin{aligned}\int_0^{1/2}\psi^2r^{N-1}\,dr&=\frac{1}{N}\psi^2r^N\Big|_{r=0}^{r=1/2}-\frac{2}{N}\int_0^{1/2}\psi\psi'r^N\,dr\\ &\le\frac{2}{N}\left(\int_0^{1/2}\psi^2r^{N-1}\,dr\right)^{1/2}\left(\int_0^{1/2}(\psi')^2r^{N+1}\,dr\right)^{1/2}.\end{aligned}$$

We apply the Young inequality $2ab\le a^2+b^2$ to infer (39).

Likewise, for (40) we have

$$\begin{aligned}
&\int_{1/2}^1 (1-r)^{\frac{p-m}{m-1}}\psi^2\,dr \\
&= \; -\frac{m-1}{p-1}(1-r)^{\frac{p-1}{m-1}}\psi^2\Big|_{r=1/2}^{r=1} + 2\frac{m-1}{p-1}\int_{1/2}^1 (1-r)^{\frac{p-1}{m-1}}\psi\psi'\,dr \\
&\leq \; 2\frac{m-1}{p-1}\left(\int_{1/2}^1 (1-r)^{\frac{2p-3}{m-1}}\psi^2\,dr\right)^{1/2}\left(\int_{1/2}^1 (1-r)^{\frac{1}{m-1}}(\psi')^2\,dr\right)^{1/2}.
\end{aligned}$$

In order to deduce (40), it remains to check that $(1-r)^{\frac{2p-3}{m-1}} \leq (1-r)^{\frac{p-m}{m-1}}$ holds if $p \geq 3-m$, and to apply the Young inequality. □

We now address Lemma 4.

Proof of Lemma 4. We derive the statement from the well-known analogous statement for regular Sobolev spaces, cf. [11, p. 272]. Let $\{\psi_n\}_{n\in\mathbb{N}}$ denote a bounded sequence in $H^1_{\rho_*}$. We may without lost of generality assume that $\int \rho_*^{p-m}\psi_n\,dx = 0$, where $p<2$ denotes a constant that satisfies the hypothesis of Lemma 3 (this is possible since $m>1$), and thus the sequence is uniformly bounded in $L^2(B_1,\rho_*^{p-m}dx)$ by (34). As $H^1_{\rho_*}$ is a separable Hilbert space, we may extract a subsequence that converges weakly to a limit ψ both in $H^1_{\rho_*}$ and in $L^2(B_1,\rho_*^{p-m}dx)$. Moreover, as ρ_* is bounded below by a positive constant in every compact subset of B_1, we have that the sequence $\{\psi_n\}_{n\in\mathbb{N}}$ is bounded in the (unweighted) Sobolev space $H^1(B_{1-\frac{1}{k}})$ for every $k\in\mathbb{N}$. By the Rellich compactness lemma, we may successively extract further subsequences $\{\psi_{\tau_k(n)}\}_{n\in\mathbb{N}}$ that converge to ψ strongly in $L^2(B_{1-\frac{1}{k}})$. By the boundedness of ρ_*, we may choose a diagonal sequence $\tau(k)=\tau_k(N(k))$ such that

$$\int_{B_{1-\frac{1}{k}}} \rho_*^{2-m}(\psi-\psi_{\tau(k)})^2\,dx \;\leq\; \frac{1}{k}, \tag{41}$$

for all $k\in\mathbb{N}$. For $\varepsilon>0$ arbitrary but fixed, we write

$$\int \rho_*^{2-m}(\psi-\psi_{\tau(k)})^2\,dx \;=\; \int_{B_{1-\varepsilon}} \rho_*^{2-m}(\psi-\psi_{\tau(k)})^2\,dx + \int_{B_1\setminus B_{1-\varepsilon}} \rho_*^{2-m}(\psi-\psi_{\tau(k)})^2\,dx.$$

The first integral converges to zero if k goes to infinity thanks to (41). For the second one, we have that

$$\int_{B_1\setminus B_{1-\varepsilon}} \rho_*^{2-m}(\psi-\psi_{\tau(k)})^2\,dx \;\lesssim\; \varepsilon^{\frac{2-p}{m-1}}\int_{B_1} \rho_*^{p-m}(\psi-\psi_{\tau(k)})^2\,dx,$$

and integrals on the right are bounded uniformly in k. Hence, letting k converge to infinity, we have that

$$\lim_{k\uparrow\infty}\int \rho_*^{2-m}(\psi-\psi_{\tau(k)})^2\,dx \;\leq\; C\varepsilon^{\frac{2-p}{m-1}},$$

for some $C>0$. Since ε was arbitrary and $p<2$, this proves the statement of Lemma 4. □

The proof of Lemma 5 is very classical and we just sketch it.

Sketch of the proof of Lemma 5. Observe that the elliptic problem is the Euler–Lagrange equation of the strictly convex functional

$$\mathcal{F}(\psi) \;=\; \frac{1}{2}\int \rho_*|\nabla\psi|^2\,dx - \int u\psi\,dx$$

for $\psi \in H^1_{\rho_*}$. Therefore, it suffices to prove existence of a minimizer of $\mathcal{F}$ in $H^1_{\rho_*}$. Indeed, by the strict convexity of $\mathcal{F}$, minimizers ψ are unique up to additive constants and satisfy the weak Euler–Lagrange equation

$$\int \rho_*\nabla\psi\cdot\nabla\xi\,dx \;=\; \int u\xi\,dx \quad \text{for all } \xi \in H^1_{\rho_*}. \tag{42}$$

In particular, choosing $\xi \in C_c^\infty(B_1)$, we see that ψ is a distributional solution of $-\nabla\cdot(\rho_*\nabla\psi) = u$, which also holds in the strong sense by interior regularity estimates. Moreover, with this information, (42) becomes

$$\int \rho_*\nabla\psi\cdot\nabla\xi\,dx \;=\; -\int \nabla\cdot(\rho_*\nabla\psi)\,\xi\,dx \quad \text{for all } \xi \in H^1_{\rho_*},$$

i.e., $\rho_*\nabla\psi\cdot\nu = 0$ on ∂B_1.

To prove existence of minimizers of $\mathcal{F}$ in $H^1_{\rho_*}$, we just mention the basic steps, following the direct method. We consider a minimizing sequence for $\mathcal{F}$ in $H^1_{\rho_*}$. With the help of Lemma 3, it is easy to check that this sequence is bounded in $H^1_{\rho_*}$. This fact is based on the estimate

$$\begin{aligned}
\int u\psi\,dx &= \int u(\psi - c)\,dx \\
&\le \left(\int \rho_*^{m-2}u^2\,dx\right)^{1/2}\left(\int \rho_*^{2-m}(\psi-c)^2\,dx\right)^{1/2} \\
&\lesssim \left(\int \rho_*^{m-2}u^2\,dx\right)^{1/2}\left(\int \rho_*|\nabla\psi|^2\,dx\right)^{1/2},
\end{aligned}$$

where c denotes the optimal constant of Lemma 3, that could be introduced in the first identity thanks to the fact that u has average zero. As a consequence, we can find a weakly converging subsequence. By lower semicontinuity of $\mathcal{F}$ with respect to weak convergence, we immediately deduce that the minimum of $\mathcal{F}$ in $H^1_{\rho_*}$ is attained. □

Acknowledgment

The author thanks Robert McCann for bringing this topic to his attention and for enlightening discussions during the process of this study. He acknowledges support through NSERC grant 217006-08.

References

[1] Abramowitz, M., and Stegun, I. A., Eds. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. Dover Publications Inc., New York, 1992. Reprint of the 1972 edition.

[2] Angenent, S. Large time asymptotics for the porous media equation. In *Nonlinear diffusion equations and their equilibrium states, I (Berkeley, CA, 1986)*, vol. 12 of *Math. Sci. Res. Inst. Publ.* Springer, New York, 1988, pp. 21–34.

[3] Barenblatt, G. I. On self-similar motions of a compressible fluid in a porous medium. *Akad. Nauk SSSR. Prikl. Mat. Meh. 16* (1952), 679–698.

[4] Beals, R., and Wong, R. *Special functions*, vol. 126 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2010. A graduate text.

[5] Benamou, J.-D., and Brenier, Y. A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. *Numer. Math. 84*, 3 (2000), 375–393.

[6] Carrillo, J. A., and Toscani, G. Asymptotic L^1-decay of solutions of the porous medium equation to self-similarity. *Indiana Univ. Math. J. 49*, 1 (2000), 113–142.

[7] Del Pino, M., and Dolbeault, J. Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions. *J. Math. Pures Appl. (9) 81*, 9 (2002), 847–875.

[8] Denzler, J., Koch, H., and McCann, R. J. Higher order time asymptotics of fast diffusion in Euclidean space (via dynamical systems methods). Preprint arXiv:1204.6434, 2012.

[9] Denzler, J., and McCann, R. J. Fast diffusion to self-similarity: complete spectrum, long-time asymptotics, and numerology. *Arch. Ration. Mech. Anal. 175*, 3 (2005), 301–342.

[10] Denzler, J., and McCann, R. J. Nonlinear diffusion from a delocalized source: affine self-similarity, time reversal, & nonradial focusing geometries. *Ann. Inst. H. Poincaré Anal. Non Linéaire 25*, 5 (2008), 865–888.

[11] Evans, L. C. *Partial differential equations*, vol. 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1998.

[12] Friedman, A., and Kamin, S. The asymptotic behavior of gas in an n-dimensional porous medium. *Trans. Amer. Math. Soc. 262*, 2 (1980), 551–563.

[13] Groemer, H. *Geometric applications of Fourier series and spherical harmonics*, vol. 61 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1996.

[14] Hebey, E. *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*, vol. 5 of *Courant Lecture Notes in Mathematics*. New York University Courant Institute of Mathematical Sciences, New York, 1999.

[15] Kamenomostskaya, S. The asymptotic behavior of the solution of the filtration equation. *Israel J. Math. 14* (1973), 76–87.

[16] Kamin, S. Similar solutions and the asymptotics of filtration equations. *Arch. Rational Mech. Anal. 60*, 2 (1975/76), 171–183.

[17] Kamin, S., and Vázquez, J. L. Fundamental solutions and asymptotic behaviour for the p-Laplacian equation. *Rev. Mat. Iberoamericana 4*, 2 (1988), 339–354.

[18] Koch, H. *Non-Euclidean singular integrals and the porous medium equation*. PhD thesis, Habilitation thesis, Universität Heidelberg, Germany, 1999.

[19] McCann, R. J. A convexity principle for interacting gases. *Adv. Math. 128*, 1 (1997), 153–179.

[20] Newman, W. I. A Lyapunov functional for the evolution of solutions to the porous medium equation to self-similarity. I. *J. Math. Phys. 25*, 10 (1984), 3120–3123.

[21] Nikiforov, A. F., and Uvarov, V. B. *Special functions of mathematical physics*. Birkhäuser Verlag, Basel, 1988. A unified introduction with applications, Translated from the Russian and with a preface by Ralph P. Boas, With a foreword by A. A. Samarskiĭ.

[22] Otto, F. The geometry of dissipative evolution equations: the porous medium equation. *Comm. Partial Differential Equations 26*, 1-2 (2001), 101–174.

[23] Otto, F., and Westdickenberg, M. Eulerian calculus for the contraction in the Wasserstein distance. *SIAM J. Math. Anal. 37*, 4 (2005), 1227–1255 (electronic).

[24] Pattle, R. E. Diffusion from an instantaneous point source with a concentration-dependent coefficient. *Quart. J. Mech. Appl. Math. 12* (1959), 407–409.

[25] Rainville, E. D. *Special functions*, first ed. Chelsea Publishing Co., Bronx, N.Y., 1971.

[26] Rainville, E. D. *Intermediate differential equations*, second ed. Chelsea Publishing Co., New York, 1972.

[27] Ralston, J. A Lyapunov functional for the evolution of solutions to the porous medium equation to self-similarity. II. *J. Math. Phys. 25*, 10 (1984), 3124–3127.

[28] Rudin, W. *Functional analysis*, second ed. International Series in Pure and Applied Mathematics. McGraw-Hill Inc., New York, 1991.

[29] Vázquez, J. L. Asymptotic beahviour for the porous medium equation posed in the whole space. *J. Evol. Equ. 3*, 1 (2003), 67–118. Dedicated to Philippe Bénilan.

[30] Vázquez, J. L. *Smoothing and decay estimates for nonlinear diffusion equations*, vol. 33 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2006. Equations of porous medium type.

[31] Vázquez, J. L. *The porous medium equation*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2007. Mathematical theory.

[32] Villani, C. *Topics in optimal transportation*, vol. 58 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003.

[33] Villani, C. *Optimal transport*, vol. 338 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2009. Old and new.

[34] Zel′dovič, Y. B., and Barenblatt, G. I. Asymptotic properties of self-preserving solutions of equations of unsteady motion of gas through porous media. *Dokl. Akad. Nauk SSSR (N.S.) 118* (1958), 671–674.

[35] Zel′dovič, Y. B., and Kompaneec, A. S. On the theory of propagation of heat with the heat conductivity depending upon the temperature. In *Collection in honor of the seventieth birthday of academician A. F. Ioffe*. Izdat. Akad. Nauk SSSR, Moscow, 1950, pp. 61–71.